

IDAHO DEPARTMENT OF FISH AND GAME

Virgil Moore, Director

Surveys and Inventories

2013 Statewide Report



MOOSE

Study I, Job 6

July 1, 2012 to June 30, 2013

Prepared by:

Wayne Wakkinen, Barb Moore Panhandle Region
George Pauley, Joshua White Clearwater Region
Randy Smith..... Magic Valley Region
Martha Wackenhut..... Southeast Region
Daryl Meints, Paul Atwood Upper Snake Region
Greg Painter, Bret Stansberry Salmon Region
Summer Crea Data Coordinator
David Smith Technical Records Specialist

Compiled and edited by: Steve Nadeau, Wildlife Staff Biologist

2013
Boise, Idaho



Findings in this report are preliminary in nature and not for publication without permission of the Director of the Idaho Department of Fish and Game.

The Idaho Department of Fish and Game adheres to all applicable state and federal laws and regulations related to discrimination on the basis of race, color, national origin, age, gender, or handicap. If you feel you have been discriminated against in any program, activity, or facility of the Idaho Department of Fish and Game, or if you desire further information, please write to: Idaho Department of Fish and Game, PO Box 25, Boise, ID 83707; or the Office of Human Resources, U.S. Fish and Wildlife Service, Department of the Interior, Washington, DC 20240.

This publication will be made available in alternative formats upon request. Please contact the Idaho Department of Fish and Game for assistance.

TABLE OF CONTENTS

STATEWIDE.....	3
PANHANDLE REGION	5
GMUS 1, 2, 3, 4, 4A, 5, 6, 7, 9.....	5
Abstract.....	5
Management Direction.....	5
Background.....	5
Population Surveys	6
Harvest Characteristics	6
Controlled Hunt Odds.....	7
Management Implications.....	7
CLEARWATER REGION	17
GMUS 8, 8A, 10, 10A, 12, 14, 15, 16, 16A, 17, 19, 20	17
Abstract.....	17
Management Direction.....	17
Population Surveys	18
Harvest Characteristics	18
Climatic Conditions	19
Management Implications.....	19
MAGIC VALLEY REGION	28
GMUS 44, 48, 49, 56, 73, 73A	28
Abstract.....	28
Management Direction.....	28
Background.....	28
Population Surveys	28
Harvest Characteristics	28
Capture and Translocation	29
Management Implications.....	29
SOUTHEAST REGION	31
GMUS 66A, 70, 71, 72, 73, 73A, 74, 75, 76, 77, 78	31
Abstract.....	31
Management Direction.....	31

TABLE OF CONTENTS (Continued)

Background31

Population Surveys31

Harvest Characteristics32

Climatic Conditions32

Habitat Conditions33

Management Implications.....33

Literature Cited33

UPPER SNAKE REGION.....42

 Abstract42

 Climatic Conditions42

 Depredation, Capture, and Translocation42

GMUS 50, 51, 58, 63, 63A45

 Background45

 Population Surveys45

 Harvest Characteristics45

 Habitat Conditions45

 Management Implications.....46

GMUS 59, 59A46

 Background46

 Population Surveys46

 Harvest Characteristics47

 Habitat Conditions47

 Management Implications.....47

GMUS 60, 60A, 61, 62, 62A47

 Background47

 Population Surveys48

 Harvest Characteristics48

 Management Implications.....49

GMUS 64, 65, 6749

 Background49

 Population Surveys49

TABLE OF CONTENTS (Continued)

Harvest Characteristics50

Habitat Conditions50

Management Implications.....50

GMUS 66, 6950

 Background.....50

 Population Surveys51

 Harvest Characteristics51

 Habitat Conditions51

 Management Implications.....51

SALMON REGION.....57

 GMUS 21, 21A, 27, 29, 30, 30A, 36A, 37A57

 Abstract.....57

 Climatic Conditions57

 Background.....57

 Population Surveys58

 Harvest Characteristics58

 Habitat Conditions58

 Capture and Translocation58

 Management Implications.....59

APPENDIX A.....63

LIST OF TABLES

PANHANDLE REGION

Table 1. Moose harvest and overall drawing odds, Panhandle Region, 1981-present.	11
Table 2. Moose harvest and drawing odds by GMU, Panhandle Region, 2000-present.	13
Table 3. Drawing odds by hunt type and season length for moose, Panhandle Region, 2005-present.....	15
Table 4. Comparison of moose harvest success rates and mean antler spread with 77-day and 14-day seasons, Panhandle Region, 2012.	16

CLEARWATER REGION

Table 1. Moose harvest and drawing odds, Clearwater Region, 1990-present.....	22
Table 2. Moose harvest and drawing odds by GMU, Clearwater Region, 2000-present.	23
Table 3. Known moose mortalities, excluding controlled hunts, Clearwater Region, 1979-present.	27

MAGIC VALLEY REGION

Table 1. Moose harvest and drawing odds by hunt area, Magic Valley Region, 1999-present.	30
---	----

SOUTHEAST REGION

Table 1. Total observed moose by sex/age class and model estimates of moose from aerial surveys, Southeast Region, 1991-2002.....	36
Table 2. Moose harvest and drawing odds, Southeast Region, 1984-present.....	37
Table 3. Moose harvest and drawing odds by hunt area, Southeast Region, 2000-present.	38
Table 4. Known moose mortalities, excluding controlled hunts, Southeast Region, 1993-present.	41

UPPER SNAKE REGION

Table 1. Moose harvest and drawing odds, Upper Snake Region, 1982-present.	52
Table 2. Known moose mortalities, excluding controlled hunts, Upper Snake Region, 1982-present.....	53
Table 3. Moose harvest and drawing odds by analysis area, Upper Snake Region, 2000-present.	54
Table 4. Aerial survey of moose, Hunt Area 62, Upper Snake Region, 2000-2001.....	56
Table 5. Aerial survey of moose, Hunt Areas 60, 60A, 61, 62, Upper Snake Region.	56

SALMON REGION

Table 1. Moose harvest and drawing odds, Salmon Region, 1990-present.	60
Table 2. Moose harvest and drawing odds by Hunt Area, Salmon Region, 2000-present.	61

LIST OF TABLES (Continued)

Table 3. Known moose mortalities, excluding controlled hunts, Salmon Region, 1982-present62

Table 4. Moose translocation, Salmon Region, February 1993.....62

LIST OF FIGURES

Figure 2. Panhandle Antlered Moose Hunt Areas.9

Figure 3. Panhandle and Clearwater Antlerless Moose Hunt Areas.....10

Figure 4. Total number of tags, antlered harvest, and antlerless harvest, Panhandle Region, 1981-2012.12

Figure 5. Number of applicants per tag, Panhandle Region, 1981-2012.12

Figure 6. Clearwater Antlered Moose Hunt Areas.20

Figure 7. Panhandle and Clearwater Antlerless Moose Hunt Areas.....21

Figure 8. Magic Valley and Southeast Antlered Moose Hunt Areas.....29

Figure 9. Southeast Antlered Moose Hunt Areas.34

Figure 10. Southeast and Upper Snake Antlerless Moose Hunt Areas.....35

Figure 11. Upper Snake and Salmon Antlered Moose Hunt Areas.43

Figure 12. Southeast and Upper Snake Antlerless Moose Hunt Areas.....44

Figure 13. Salmon Antlered Moose Hunt Areas.....59

STATEWIDE REPORT SURVEYS AND INVENTORY

JOB TITLE: Moose Surveys and Inventories

STUDY NAME: Big Game Population Status, Trends, Use, and Associated Habitat Studies

PERIOD COVERED: July 1, 2012 to June 30, 2013

STATEWIDE

Moose have expanded their range and numbers in parts of Idaho over the past few decades, moving westward into Washington and northeastern Oregon and southward into Utah. Harvest records and hunter reports indicate however, that moose populations in central Idaho Wilderness and other areas of the Clearwater continue to decline. A variety of reasons may be to blame and research is being conducted to identify the causes. Also, across the state moose permits have fluctuated in response to reduced success rate, reduced antler size, and reduced numbers of animals seen by hunters. Moose range has increased in some areas but numbers have declined in others in every region across the state. Statewide harvest has declined since the peak in the mid 2000's.

A total of 554 antlered moose were reported harvested by 741 tag holders in fall 2012 (Table 1). The mean antler spread of harvested moose was 37.7 inches (up from 36 inches in 2011), based on animals measured during the mandatory check conducted statewide at Regional Offices, taxidermists, and contracted checkpoints. Based on 696 reports received (no reports were received from 45 tag holders), harvest success on antlered moose averaged over 80% statewide.

In addition, 120 antlerless moose (including 3 antlerless males) were harvested by the 154 tag holders in fall 2012 (Table 1). The hunter success rate of antlerless moose based on 145 reports received was 83%.

An additional 4 tags were issued in conjunction with the Department's "Super Tag" drawings. Four antlered moose (100%, up from 75% in 2011) were reported harvested, in Units 1, 55, 56 and 67. These 4 moose had an average antler spread of 40.4 inches.

Data on moose age and antler spread at harvest were analyzed to assist in the monitoring and evaluation of current and creation of new hunting seasons.

Moose continue to be one of Idaho's most desirable trophy species among resident hunters. Hunters are allowed to draw a tag to harvest only 1 antlered and 1 antlerless moose in their lifetime (except for those tags left over after the initial drawing, which do not apply to the lifetime limit). A total of 5,637 first-choice applications were received for the 695 tags for antlered moose in April 2013 for the fall 2013 hunting season, yielding overall drawing success of 12%. No antlered tags were available after the first drawing. Most applicants for antlered moose tags were resident Idahoans (5,190 or 92% of the total); only 447 non-residents applied despite non-residents being able to draw up to 10% of the total number of tags offered.

Of the 870 applicants for 164 antlerless moose tags allocated among 25 different hunt areas, 866 (99.5%) were received from residents. No antlerless tags were available after the first drawing.

Table 1. Moose hunter participation and harvest between July 1, 2012 and June 30, 2013.

Area	Hunters	Hunter Days	Total Harvest	Males	Females	% Change in Total Harvest from Previous Year
Statewide	899	4,009	678	558	120	-3%

PANHANDLE REGION

GMUs 1, 2, 3, 4, 4A, 5, 6, 7, 9

Controlled Hunt Areas 1-1, 1-2, 1-3, 1-4, 2, 3, 4, 4A, 5, 6, 7, 9

Abstract

The Panhandle Region offered 358 tags in the 2012 season, over three times the number of tags offered ten years ago. However, this represents a reduction of 15 tags from 2011 because of possible declining populations or hunt quality in some controlled hunt areas. Overall drawing odds for moose tags in the region was 12.4%, consistent with the previous 10 years and substantially better than prior odds. During 2012, 8 of the 261 harvested bulls for which we have antler spread measurements were at least 50 inches in width (3.1%). The average spread for harvested antlered moose (n=261) was 37.4 inches. Success rates averaged 83% from 2002-2011 and was 82% in 2012.

Management Direction

1. Develop an index to moose population trends that does not rely solely on aerial surveys.
2. Place enforcement emphasis on known problem areas of illegal moose kills. Publicize moose poaching arrests and the statewide reward system (Citizens Against Poaching) in the media.
3. Develop a program for warning deer and elk hunters that moose are in an area to reduce accidental kills of moose.
4. Examine present controlled hunt boundaries and permit levels and adjust as necessary to provide quality hunts and distribute moose hunters more evenly. Coordinate moose management and tag levels along the Idaho/Washington border with the Washington Department of Fish and Wildlife.
5. Continue collecting information on moose distribution and mortality from Department and other agency personnel and the hunting public.

Background

Open areas and extensive riparian areas that typify moose habitat elsewhere are not widespread in the Panhandle Region. Moose in this region often utilize closed-canopy timber stands with interspersed shrub fields and creek bottoms. Presently, moose populations appear to be stable in most areas of the Panhandle (Figures 1 & 2).

Historically, moose have been managed in Idaho for long hunts with high success rates and a good opportunity to harvest a large-antlered bull. This conservative approach, coupled with a high demand for moose hunting, led to poor odds for drawing a moose tag. In response, short, 7-day hunts were initiated during the fall of 2005 to: a) provide hunters a choice for better drawing odds at the expense of season length and b) provide data on how success rates change with a short season. Further modifications to the moose hunting season structure were initiated for the 2007 and 2008 seasons. The 86-day hunts in Game Management Units (GMUs) 1 and 2 were eliminated and replaced with a series of 14-day hunts.

For the 2009-2010 and 2011-2012 seasons, both long and short hunts were offered. Long seasons (77 days) offered more opportunity but lower drawing odds while short hunts resulted in better drawing odds. Hunters applying for the short hunts were over twice as likely to successfully draw a tag (Table 3).

Population Surveys

In December of 2010, an aerial helicopter survey was flown in the northern part of GMU 5, including the Mica Peak area from the Spokane River south to Windy Bay. In 18 search units, 68 total moose were observed (23 bulls, 26 cows, 15 calves, 4 un-classified). The moose sightability data were run through several models. Quayle et al. (2001) developed a sightability model for moose in south-central British Columbia, which estimated the Mica Peak population to be 72. Anderson and Lindzey's (1996) sightability model developed for moose in Wyoming estimated the total to be 82 moose. The Hiller sightability model, developed for elk, produced an estimate of 100 moose. The Bell helicopter model, developed for elk, estimated the total number of moose in the Mica Peak area to be 115.

An aerial thermal infrared survey was conducted by Vision Air Research, Boise, ID at the end of March in the same section of GMU 5. Transects were flown 800 ft. apart at 1,500 ft. above ground in a fixed wing aircraft. An infrared sensor was mounted on the wing of the aircraft and operated by a wildlife biologist. Moose were located by observing their level of emitted infrared energy versus background levels. A total of 112 moose were observed. Some moose may have been missed if they were obscured by vegetation.

The infrared survey technique yielded an estimate for the number of moose that fell within the range of the elk sightability model estimates. Thermal infrared surveys should be further explored as a way to estimate moose populations.

Harvest Characteristics

Moose hunting was authorized in all Panhandle GMUs for the first time in 2007 (Table 2). In 2007, 5 antlered tags each were issued in GMU 4A and GMU 5. The Department issued 358 moose tags for the 2012 season: 195 tags for antlered moose with a 77-day season (15 Sept – 1 Dec), 118 tags for antlered moose with 2 different 14-day seasons (1 Oct – 14 Oct; 1 Nov – 14 Nov), and 45 tags for antlerless moose with a 48-day season (15 Oct – 1 Dec).

Hunters reported harvesting 305 moose with the 358 available tags for an overall success rate in 2011 of 85% (Table 1). This is the same as the average success rate over the past 20 years of moose harvest throughout the Panhandle. Success rates in individual GMUs varied from 40% to 100%, but small sample sizes in some of these GMUs make success rates volatile.

In 2012, 8 of the 261 harvested bulls for which we have antler spread measurements were equal to or exceeded 50 inches (3.1%). This is similar to previous data for the previous 5 years and is as follows: 2011: 2.0%, 2010: 2.8%, 2009: 1.0%, 2008: 0.8%, 2007: 2.3%, and 2006: 3.7%.

Controlled Hunt Odds

Most areas of Idaho have tags available for a variety of big game species. By forcing a choice between moose and other big game tags, the Department has been successful in substantially improving drawing odds across most of the state. In the Panhandle, the only big game species managed entirely under a tag system is moose, making drawing odds poor for moose.

In an attempt to address the complaint of hunters that it was too difficult to draw a moose tag, the Department conducted a trial 7-day hunt for 2005 and 2006 to provide an avenue for improving drawing odds. It was believed that relatively few hunters would opt for the shorter season, thus greatly improving drawing odds for those hunters who were interested in choosing better drawing odds at the expense of a shorter hunting season. It was also believed that success rates would diminish slightly with the shorter season, allowing the moose herd to support additional tags to be issued, which would further improve drawing odds.

Over the past 29 years, the number of moose applicants in the Panhandle Region has steadily risen, but the number of tags being offered has increased at a faster rate, resulting in significantly better drawing odds (Table 1, Figures 3 & 4). Further, antlered moose hunts with short seasons had much better drawing odds than longer seasons (Table 3).

Another modification of the shorter hunts was offered in 2007-2008. A series of 6 14-day hunts were offered in GMUs 1 and 2 with the first hunt starting on 30 August and the last hunt starting on 15 November. This was another attempt to provide hunter opportunity and improve drawing odds. Drawing odds were significantly better for these 14-day hunts as compared to the traditional 86-day hunts, however, hunters were disappointed that no long hunt was offered in these GMUs.

For the 2009-2010 and 2011-2012 season, both long and short hunts were offered in most Panhandle GMUs. Drawing odds were better for the shorter hunts (Table 3) and hunters seemed happy that they had the choice of the long or short hunts.

Management Implications

An attempt was made beginning in 2001 to become less conservative in many of our moose hunts, particularly in Hunt Areas 1-1, 1-3, and 2. The overall drawing odds have improved to the point that an applicant now has a one in 8 chance of drawing a moose tag in the Panhandle Region. Success rates have remained relatively high and the mean antler spread has remained stable.

The lack of moose population surveys is a serious handicap to moose management in Idaho. For the most part, tag levels continue to be set conservatively, based on anecdotal information and the perception of what is socially acceptable. This conservative approach has produced poor drawing odds, the major complaint regarding moose management in Idaho, although recent changes in the Panhandle Region have improved the situation. However, the lack of surveys makes it difficult to determine the impact of the significant changes that have been made to the Panhandle seasons.

Drawing odds were much better for the 14-day hunts than the 77-day hunts, providing an avenue for hunters willing to trade season length for improved odds. Hunters with the shorter hunts reported high satisfaction with the hunts during animal check-ins. It was hypothesized that the success rates for the shorter hunts would be lower than the longer hunts, allowing more hunters afield. The difference, however, was relatively minor. The success rates during the different time periods of these short hunts will be used to evaluate the practicality of continuing to offer these hunts and the possibility of adjusting tag levels based on success rates.




Figure 1. Panhandle Antlered Moose Hunt Areas.




Figure 2. Panhandle and Clearwater Antlerless Moose Hunt Areas.

Table 1. Moose harvest and overall drawing odds, Panhandle Region, 1981-present.

Year	Tags	Harvest			Hunter success (%)	First-choice applicants	Drawing Odds (%)
		M	F	Total			
1981	11	7	0	7	64	701	2
1982	11	11	0	11	100	599	2
1983	15	14	0	14	93	712	2
1984	15	14	0	14	93	721	2
1985	28	21	0	21	75	907	3
1986	28	23	0	23	82	750	4
1987	28	24	0	24	86	653	4
1988	40	34	0	34	85	597	7
1989	40	35	0	35	88	725	6
1990	42	38	0	38	90	849	5
1991	51	45	0	45	88	1,024	5
1992	51	44	0	44	86	1,071	5
1993	83	69	0	69	83	1,361	6
1994	83	63	0	63	76	1,430	6
1995	100	84	0	84	84	1,529	7
1996	100	74	0	74	74	1,516	7
1997	103	85	0	85	83	1,837	6
1998	103	91	0	91	88	1,623	6
1999	123	100	0	100	81	2,001	6
2000	123	106	0	106	86	1,765	7
2001	220	176	5	181	82	1,799	12
2002	220	156	5	161	73	1,703	13
2003	235	189	17	206	88	1,858	13
2004	236	188	14	202	86	2,088	11
2005	285	226	26	253	88	2,536	11
2006	285	215	22	237	83	2,878	10
2007	352	251	32	283	80	2,443	14
2008	352	235	36	271	77	2,352	15
2009	386	298	48	346	90	2,763	14
2010	386	283	50	333	86	2,814	14
2011	388	277	46	323	83	3,136	12
2012 ^a	358	263	42	305	85	2,880	12

^a No tags (15) were sold in hunt numbers 3009 and 3095 in 2012.




Figure 3. Total number of tags, antlered harvest, and antlerless harvest, Panhandle Region, 1981-2012.




Figure 4. Number of applicants per tag, Panhandle Region, 1981-2012.

Table 2. Moose harvest and drawing odds by GMU, Panhandle Region, 2000-present.

GMU	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing Odds (%)
			M	F				
1	2000	88	75	0	85	8.6	812	11
	2001	155	120	0	77	8.6	828	19
	2002	155	103	0	66	9.2	1,065	15
	2003	170	135	14	88	9.3	1,165	15
	2004	171	131	10	82	7.2	1,185	14
	2005	170	145	18	96	8.9	1,220	14
	2006	170	139	15	90	8.1	1,316	13
	2007	218	147	17	75	8.7	1,053	21
	2008	218	136 ^a	18	71	5.6	917	24
	2009	206	160 ^a	15	85	7.0	1,112	19
	2010	206	154	20	84	5.7	1,071	19
	2011	180	131	7	77	7.9	1,056	17
2012 ^{a,b}	151	119	1	79	8.1	809	18	
2	2000	10	10	0	100	6.4	162	6
	2001	25	20	5	100	7.1	211	12
	2002	25	20	5	100	4.4	205	12
	2003	25	20	4	96	8.2	208	12
	2004	25	17	4	84	5.5	287	9
	2005	35	25	8	94	6.0	309	11
	2006	35	25	7	91	6.5	385	9
	2007	44	25	15	91	6.9	334	13
	2008	44	22	18	91	2.8	496	9
	2009	65	35	28	97	5.6	526	12
	2010	65	31	25	86	7.5	506	13
	2011	75	10	30	93	4.8	673	11
	2012	70	36	33	99	4.4	746	9
3	2000	5	4	0	80	11.3	27	19
	2001	5	5	0	100	7.2	35	14
	2002	5	5	0	100	10.8	49	10
	2003	5	4	0	80	8.5	44	11
	2004	5	5	0	100	6.8	66	8
	2005	10	11 ^a	0	100	4.9	83	12
	2006	10	10	0	100	3.9	114	9
	2007	20	19	0	95	7.2	122	16
	2008	20	18	0	90	5.9	165	12
	2009	30	24	5	97	5.6	192	16
	2010	30	20	5	83	6.2	227	13
	2011	33	25	4	88	5.2	230	14
	2012	32	26	4	94	5.0	259	12
4	2000	5	5	0	100	9.5	68	7
	2001	10	9	0	90	12.0	108	9
	2002	10	7	0	70	10.0	122	8
	2003	10	8	0	80	14.6	133	8
	2004	10	8	0	80	9.9	175	6
	2005	15	15	0	100	4.0	229	7
	2006	15	13	0	87	8.1	247	6
	2007	20	20	0	100	8.2	333	6

Table 2 Continued

GMU	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing Odds (%)
			M	F				
	2008	20	19	0	95	4.4	364	5
	2009	25	22	0	88	9.7	358	7
	2010	25	25	0	100	5.0	398	6
	2011	30	28	0	93	7.0	471	6
	2012	30	28	0	93	6.86	386	8
4A	2007	5	2	0	40	3.0	20	25
	2008	5	2	0	40	12.5	24	21
	2009	5	4	0	80	3.0	8	63
	2010	5	4	0	80	3.0	17	29
	2011	5	2	0	40	12.5	13	39
	2012	5	2	0	40	16.5	14	36
5	2007	5	5	0	100	7.3	163	3
	2008	5	4	0	80	9.3	149	3
	2009	10	11 ^a	0	100	6.8	175	6
	2010	10	10	0	100	11.9	193	5
	2011	20	12	5	85	5.9	217	9
	2012	20	14	4	90	5.3	249	8
6	2000	5	4	0	80	8.3	121	4
	2001	10	7	0	70	11.0	132	8
	2002	10	8	0	80	4.1	147	7
	2003	10	10	0	100	9.2	185	5
	2004	10	8	0	80	9.9	233	4
	2005	15	14	0	93	6.4	275	5
	2006	15	13	0	87	6.9	334	4
	2007	20	20	0	100	7.2	292	7
	2008	20	20 ^a	0	100	5.8	338	6
	2009	25	26 ^a	0	100	6.7	294	9
	2010	25	24	0	96	7.1	280	9
	2011	25	23	0	92	6.1	321	8
	2012	25	23	0	92	6.57	289	9
7	2000	5	3	0	60	8.8	34	15
	2001	10	10	0	100	11.8	108	9
	2002	10	10	0	100	9.4	57	18
	2003	10	9	0	90	5.0	83	12
	2004	10	8	0	80	4.1	86	12
	2005	10	8	0	80	4.7	112	9
	2006	10	7	0	70	12.0	97	10
	2007	10	9	0	90	6.9	70	14
	2008	10	5	0	50	6.8	68	15
	2009	10	9	0	90	4.4	36	28
	2010	10	8	0	80	4.9	68	15
	2011	10	8	0	80	6.0	51	20
	2012	10	7	0	70	4.0	51	20
9	2000	5	5	0	100	9.2	41	12
	2001	5	5	0	100	8.0	61	8
	2002	5	5	0	100	10.0	40	13
	2003	5	5	0	100	10.8	40	13
	2004	5	5	0	100	8.0	56	9

Table 2 Continued

GMU	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing Odds (%)
			M	F				
	2005	10	9	0	90	5.8	54	19
	2006	10	8	0	80	4.4	69	14
	2007	10	9	0	90	6.9	56	18
	2008	10	9	0	90	6.4	78	13
	2009	10	9	0	90	4.1	62	16
	2010	10	10	0	100	4.9	54	19
	2011	10	8	0	80	5.3	56	18
	2012	10	8	0	80	4.2	48	21

^a Includes one Supertag harvest.

^b No tags (15) were sold in hunt numbers 3009 and 3095 in 2012.

Table 3. Drawing odds by hunt type and season length for moose, Panhandle Region, 2005-present.

Year	Hunt type	Season length (days)	Tags	Applicants per tag
2005	Antlered	86	200	11.0
	Antlered	7	55	1.5
	Antlerless	40	30	8.5
2006	Antlered	86	200	12.0
	Antlered	7	55	4.6
	Antlerless	40	30	7.2
2007	Antlered	86	50	18.5
	Antlered	14	262	4.8
	Antlerless	40	40	6.7
2008	Antlered	86	50	18.3
	Antlered	14	262	4.6
	Antlerless	40	40	6.2
2009	Antlered	77	210	9.4
	Antlered	14	121	3.3
	Antlerless	49	55	7.3
2010	Antlered	77	210	9.2
	Antlered	14	121	4.0
	Antlerless	49	55	7.3
2011	Antlered	77	205	10.5
	Antlered	14	128	4.0
	Antlerless	48	55	9.2
2012	Antlered	77	195	9.7
	Antlered	14	118	4.7
	Antlerless	48	45	7.7

Table 4. Comparison of moose harvest success rates and mean antler spread with 77-day and 14-day seasons, Panhandle Region, 2012.

Season length	Season Dates	Tags issued	Number harvest	Success rate (%)	Mean antler spread
77 days	15 Sep –1 Dec	205	167	81	36.9
14 days	1 Oct – 14 Oct	60	46	77	38.3
14 days	1 Nov – 14 Nov	68	50	74	36.2

CLEARWATER REGION

GMUs 8, 8A, 10, 10A, 12, 14, 15, 16, 16A, 17, 19, 20

**Controlled Hunt Areas 8, 8A, 10-1, 10-2, 10-3, 10-4, 10-5, 10-6,
10A-1, 10A-2, 10A-3, 10A-4, 10A-5, 12-1, 12-2, 12-3, 12-4, 12-5, 12-6,
14-1, 14-2, 15-1, 15-2, 15-3, 15-4, 16-1, 16-2, 16A, 17, 19, 20**

Abstract

Based upon mandatory harvest report data, Clearwater Region hunters harvested 64 antlered moose in 28 antlered-only controlled hunts and an additional 6 antlerless moose in 2 controlled hunts for antlerless moose in 2012. A total of 6 antlered-only permit holders opted to take rain checks in Unit 14 out of 20 due to severely limited access as result of large wildfires. The 147 hunters in the region harvested a total of 70 moose. A total of 153 (145 antlered, 8 antlerless) tags were available across the region for a total harvest success rate of 48%. Antlered and antlerless success rates were 46% and 75%, respectively. Drawing odds ranged from 1:1.0 (Hunt Areas 10-4, 10-6, 12-3, 12-4, 12-6, 16-2, 16A, and 20) to 1:21.4 (Hunt Area 8A). The mean antler spread for the 64 antlered moose harvested in the region was 38.5 inches, with a range of 16 to 55 inches for 2012. Cumulative drawing odds for antlered-only hunts in the Clearwater Region were 1:4.2 for the 2012 season.

Management Direction

Moose populations will be allowed to increase in GMUs where habitat conditions will support expansion. Legal harvest will continue to be focused on antlered bulls. Antlerless moose hunting opportunities will be continued in those areas where population control measures are considered desirable. Moose harvest will be increased where feasible and decreased where necessary. Known mortalities will be documented and information on numbers and distribution will be obtained from big game mandatory harvest checks.

Moose populations large enough to support hunts are found in most of the Clearwater except GMUs 11, 11A, 13, and 18. GMUs are divided into controlled hunts to disperse hunters and to direct harvest to specific areas (Figures 5 & 6).

Historically, moose were hunted through controlled hunts on a bulls-only basis; however, in 1999, 2 antlerless moose hunts (Hunts 8-2 and 8A-2 with 4 tags each) were initiated to increase hunting opportunity, address high cow moose densities, and minimize the potential for moose-automobile collisions in these areas. Hunting season lengths for moose in the Clearwater Region were 86 days for antlered moose hunts and 40 days for antlerless hunts (Appendix A). Since 1986, persons applying for moose tags have been prohibited from applying for any other controlled hunt to improve drawing odds. Additionally, unsuccessful tag holders must wait 2 years before applying for another controlled moose hunt. Tag levels are based on trends in antler spread of harvested moose and hunter success rates of recent tag holders in the respective controlled hunts.

Some moose populations in the Clearwater Region are found in climax vegetative cover. Summer feeding habits tend to be nocturnal in open, wet meadows, while diurnal activity is

limited to adjacent forested areas. Logging may improve or reduce habitat for these populations. Winter habitat selection favors young subalpine fir and Pacific yew plant communities. Other populations are adapted to early seral plant communities, except in winter. These populations may be expanding in areas where extensive habitat manipulation has resulted in seral brush fields. Winter ranges appear to be timbered areas where yew-wood thickets are several hundred years old. Creating openings in these timber stands through logging may impact moose by eliminating these yew-wood thickets. Effects of the recent expansion of wolves on moose populations within the region are as yet undetermined.

Population Surveys

Moose in the Clearwater Region are usually counted incidental to elk surveys. Consequently, many moose are not counted because these surveys are seldom flown at elevations where moose normally winter and because moose tend to prefer dense subalpine fir plant associations for winter habitat where they are less conspicuous. As a result, no comparative population data have been collected on a regular basis on moose throughout the region.

A sightability survey of moose in GMU 15 was attempted in 2000. Results were unsatisfactory because of overly large confidence intervals. These results were due to the extreme correction factors applied to animals detected under heavy canopy coverage classes. During model development, only 4 moose were encountered and in cover greater than 70%.

Harvest Characteristics

Harvest levels, hunter success, and hunter days expended for 2011 were determined from mandatory harvest reports (Tables 1 and 2). Several changes have been made to regional moose Hunt Areas in recent history. Hunt areas in GMUs 12, 15, and 17 were combined and/or renamed in 2001 and 1 new hunt area was added in GMU 10 (10-6) in 2001. Tag numbers were adjusted in the region to respond to changes in hunter success rates and/or antler spread with a net loss of 22 tags in 2001 (from 290 to 270) and a further reduction of 20 tags in 2005 (250). For the 2009 and 2010 seasons, the total number of tags was reduced from 250 (242 antlered and 8 antlerless) to 169 (161 antlered and 8 antlerless). A large portion of this reduction came about as result of a reconfiguration (elimination of numerous small hunts into one unit-wide hunt with reduced tag levels in Units 16A, 17, 19 and 20). Antlered-only tags were further reduced in 2011 from 161 to 145 as result of combining the 4 hunts (with 24 tags) in GMU 15 into 1 GMU-wide hunt with 8 tags. The 153 moose tags available in 2012 resulted in a reported harvest of 64 antlered and 6 antlerless moose.

The 2012 cumulative success rate of 48% was lower than the previous 5-year (2007-2011) average of 51%. Success rates for 2012 antlered and antlerless moose were 46% and 75%, respectively. Drawing odds ranged from 1:1.0 (Hunt Areas 10-4, 10-6, 123, 12-4, 12-6, 16-2, 16A and 20) to 1:21.4 (Hunt Area 8A).

Reported moose mortalities due to methods other than legal harvest during controlled hunts have varied considerably by year (Table 3). It is likely that the level of mortality is considerably higher than what is reported.

The mean antler spread for the 64 antlered moose harvested in the region in 2012 was 38.5 inches, with a range of 16 to 55 inches. Cumulative drawing odds for antlered-only hunts in the Clearwater Region were 1:4.2 in 2012.

Climatic Conditions

According to the United States Department of Agriculture Natural Resources Conservation Service, heavy precipitation was recorded in October of 2012, with several sites receiving twice their normal amount. January 2013 produced 84% of normal monthly precipitation keeping snowpack percentages similar to the previous month and bringing the precipitation level to 107% of average since October 1, 2012; below normal precipitation also fell during February and March, 75% and 82% of average, respectively. This dry trend continued through May, however the Clearwater basin's water supply was better than any Idaho basin to its south. Snow amounts were 81% of the 1981-2010 median as of June 1, 2013. By watershed, the North Fork Clearwater, Lochsa, and Selway Rivers were 89%, 49%, and 41% of normal snowpack, respectively, as of 1 June. Compared to last year's 1 June snowpack of 109% of average, this year's reduced snowpack may have contributed to improved survival of deer and elk in backcountry areas.

Management Implications

Tag levels will continue to be allocated based on trends in antler spread of harvested moose and hunter success rates of recent tag holders. Numbers of tags may be increased or decreased as dictated by harvest data. Tag numbers were decreased by 22 in the Clearwater Region in 2001 and by an additional 20 tags in 2005. Tag numbers were again reduced for the 2009 season by an additional 81 tags.

All areas need more intensive investigation to determine population levels, trends, and habitat selection and use. Some moose populations appear to be increasing and seem to respond favorably to extensive habitat alteration by silvicultural practices. However, other populations may be displaced or eliminated because they cannot adapt to habitat changes, particularly where yew-wood thickets are eliminated through logging and where increased road densities make moose more vulnerable to illegal and Native American harvest.

Additionally, the effects of the recent expansion of wolves across the region on moose populations are as yet undetermined. In 2008, the region began monitoring moose in GMU 10 that were captured and radio-collared to determine mortality rates and causes of death in the presence of wolves. This work is being done in conjunction with the ongoing wolf-elk interaction research in the Lolo Zone. A total of 12 radio-collars were placed on yearling or adult moose during the 2008-2009 winter. Eleven of the 12 collared animals survived the first year. The lone mortality was a young bull that was harvested by a hunter in Hunt Area 10-3 in 2009. One additional radio-collar was deployed in January 2010 prior to a helicopter accident, after which capture operations ceased. Again, 11 of the 12 collared animals survived the year (2010). The one mortality was a bull that was injured while sparring with another bull during the rut. In February 2011, an additional 22 moose were captured and radio-collared (2 bulls, 8 cows, and 12 calves). Since January 2012, wolves had killed 1 adult cow moose and 6 calves in addition to 2 unknown cow and 1 non-predation bull mortalities. There were no additional

collars deployed in 2012. In 2013, an additional 3 cow and 3 calf moose were captured and fitted with radio-collars. While results are very preliminary, to date, wolves have not proven to be a significant cause of mortality on radio-collared adult moose. However, if early trends in wolf-caused calf mortality continue, calf survival and recruitment could be a serious issue.




Figure 5. Clearwater Antlered Moose Hunt Areas.




Figure 6. Panhandle and Clearwater Antlerless Moose Hunt Areas.

Table 1. Moose harvest and drawing odds, Clearwater Region, 1990-present.

Year	Tags	Harvest			Hunter success (%)	First-choice applicants	Drawing odds
		M	F	Total			
1990	167	118	0	118	71	1,156	1:6.9
1991	176	134	0	134	76	1,201	1:6.8
1992	176	132	0	132	75	1,221	1:6.9
1993	201	159	0	159	79	1,211	1:6.0
1994	201	133	0	133	66	1,115	1:5.5
1995	263	177	0	177	67	1,501	1:5.7
1996	263	162	0	162	62	1,288	1:4.9
1997	263	157	0	157	60	1,579	1:6.0
1998	263	153	0	153	58	1,250	1:4.8
1999	292	180	8	188	64	1,540	1:5.3
2000	292	177	7	184	63	961	1:3.3
2001	270	141	7	148	55	931	1:3.4
2002	270	151	8	159	59	813	1:3.0
2003	270	156	6	162	60	798	1:3.0
2004	270	150	7	157	58	891	1:3.3
2005	250	152	8	160	64	964	1:3.9
2006	250	144	7	151	60	943	1:3.8
2007	250	130	7	137	55	938	1:3.8
2008	250	117	8	125	50	850	1:3.4
2009	169	79	6	85	50	788	1:4.7
2010	169	79	8	87	51	801	1:4.5
2011	156	71	7	78	51	625	1:4:1
2012	147 ^a	64	6	70	48	644	1:4.2

^a 153 permits were issued in 2012, although 6 hunters elected to take rain checks (wildfire related) and did not participate in the hunt.

Table 2. Moose harvest and drawing odds by GMU, Clearwater Region, 2000-present.

Hunt area	Year	Tags	Harvest		Hunter success (%)	Days/hunter ^a	First-choice applicants	Drawing odds
			M	F				
8	2000	10	5	3	80	5.1	34	1:3.4
	2001	10	5	3	80	7.1	35	1:3.5
	2002	10	6	4	100	5.4	52	1:5.2
	2003	10	6	3	90	5.4	48	1:4.8
	2004	10	6	4	100	4.2	54	1:5.4
	2005	12	8	4	100	12.0	66	1:5.5
	2006	12	7	4	92	8.3	73	1:6.1
	2007	12	7	4	92	6.5	98	1:8.2
	2008	12	7	4	92	3.1	112	1:9.3
	2009	12	7	4	92	5.3	123	1:10.3
	2010	12	7	4	92	3.0	164	1:13.7
	2011	12	8	4	100	7.9	144	1:12.0
2012	12	7	3	83	4.3	144	1:12.0	
8A	2000	10	6	4	100	3.5	76	1:7.6
	2001	10	5	4	90	4.1	104	1:10.4
	2002	10	5	4	90	4.6	93	1:9.3
	2003	10	6	3	90	11.3	113	1:11.3
	2004	10	6	4	100	6.8	105	1:10.5
	2005	12	8	4	100	8.2	138	1:11.5
	2006	12	7	3	83	10.4	142	1:11.8
	2007	12	8	3	92	7.7	169	1:14.1
	2008	12	8	4	100	6.5	181	1:15.1
	2009	12	8	2	83	7.9	201	1:16.8
	2010	12	8	4	100	7.5	223	1:18.6
	2011	12	8	3	92	5.5	171	1:14.3
2012	12	8	3	92	10.4	218	1:18.2	
10	2000	23	13	0	57	4.0	112	1:4.9
	2001	28	17	0	61	6.4	91	1:3.3
	2002	28	14	0	50	9.3	86	1:3.1
	2003	28	20	0	71	6.4	82	1:2.9
	2004	28	21	0	75	3.9	105	1:3.8
	2005	32	21	0	66	7.8	100	1:3.1
	2006	32	20	0	63	9.2	112	1:3.5
	2007	32	25	0	78	5.7	113	1:3.5
	2008	32	17	0	53	6.6	106	1:3.3
	2009	32	22	0	69	9.2	120	1:3.8
	2010	32	19	0	59	5.8	97	1:3.0
	2011	32	13	0	41	6.2	77	1:2.4
2012	32	14	0	44	2.7	48	1:15	
10A	2000	34	29	0	85	11.9	134	1:3.9
	2001	32	28	0	88	6.8	116	1:3.6
	2002	32	26	0	81	7.9	130	1:4.1
	2003	32	27	0	84	8.9	140	1:4.4
	2004	32	25	0	78	9.4	145	1:4.5

Table 2. Continued.

Hunt area	Year	Tags	Harvest		Hunter success (%)	Days/hunter ^a	First-choice applicants	Drawing odds
			M	F				
12	2005	34	32	0	94	7.6	148	1:4.4
	2006	34	26	0	76	7.6	172	1:5.1
	2007	34	31	0	91	11.8	191	1:5.6
	2008	34	24	0	71	9.0	192	1:5.6
	2009	29	20	0	69	13.9	168	1:5.8
	2010	29	20	0	69	6.1	152	1:5.2
	2011	29	19	0	66	8.2	131	1:4.5
	2012	29	16	0	55	5.4	115	1:4.0
	2000 ^b	61	31	0	51	6.3	119	1:2.0
	2001	45	16	0	36	3.0	70	1:1.6
	2002	45	24	0	53	4.5	58	1:1.3
	2003	45	27	0	58	6.7	75	1:1.7
	2004	45	22	0	49	5.6	87	1:1.9
	2005	43	20	0	47	6.9	73	1:1.7
	2006	43	23	0	53	8.5	70	1:1.6
	2007	43	18	0	42	9.0	73	1:1.7
	2008	43	21	0	49	10.6	64	1:1.5
	2009	26	9	0	35	5.9	42	1:1.6
	2010	26	15	0	58	11.1	48	1:1.8
	2011	26	10	0	38	5.9	27	1:1.1
2012	26	13	0	50	3.2	44	1:1.7	
14	2000	10	9	0	90	4.5	100	1:10.0
	2001	13	11	0	85	3.5	124	1:9.5
	2002	13	11	0	85	5.3	120	1:9.2
	2003	13	11	0	85	4.6	121	1:9.3
	2004	13	11	0	85	8.2	114	1:8.8
	2005	13	11	0	85	10.0	114	1:8.8
	2006	13	10	0	77	10.4	92	1:7.1
	2007	13	8	0	62	6.5	71	1:5.5
	2008	13	6	0	46	8.0	83	1:6.4
	2009	11	6	0	56	9.2	42	1:3.8
	2010	11	5	0	45	4.4	55	1:5.0
	2011	11	6	0	56	12.8	33	1:3.0
15	2012	7 ^c	2	0	29	0.5	37	1:3.4
	2000	60	44	0	73	8.2	212	1:3.5
	2001	60	34	0	57	8.9	256	1:4.3
	2002	60	35	0	58	8.5	176	1:2.9
	2003	60	35	0	58	11.2	173	1:2.9
	2004	60	37	0	62	7.1	186	1:3.1
	2005	45	30	0	67	8.4	155	1:3.4
	2006	45	25	0	55	12.4	143	1:3.2
	2007	45	20	0	44	11.1	117	1:2.6
	2008	45	18	0	40	11.0	108	1:2.4
	2009	24	3	0	13	6.0	70	1:2.9

Table 2. Continued.

Hunt area	Year	Tags	Harvest		Hunter success (%)	Days/hunter ^a	First-choice applicants	Drawing odds
			M	F				
16	2010	24	4	0	17	25.8	46	1:1.9
	2011	8	2	0	25	12.0	27	1:3.4
	2012	8	1	0	13	0.3	13	1:1.6
	2000	14	13	0	93	6.2	78	1:5.6
	2001	17	10	0	59	6.3	65	1:3.8
	2002	17	11	0	65	5.4	40	1:2.4
	2003	17	9	0	53	7.0	58	1:3.4
	2004	17	10	0	59	4.8	47	1:2.8
	2005	12	8	0	67	6.3	55	1:4.6
	2006	12	6	0	50	5.7	37	1:3.1
	2007	12	9	0	75	8.2	38	1:3.2
	2008	12	3	0	25	12.7	38	1:3.2
	2009	4	2	0	50	6.5	7	1:1.8
	2010	4	1	0	25	ND	7	1:1.8
16A	2011	4	2	0	50	6.8	5	1:1.3
	2012	4	0	0	0	ND	4	1:1.0
	2000	7	3	0	43	8.7	21	1:3.0
	2001	7	6	0	86	4.3	13	1:1.9
	2002	7	3	0	43	14.3	14	1:2.0
	2003	7	3	0	43	4.0	8	1:1.1
	2004	7	5	0	71	16.8	12	1:1.7
	2005	7	5	0	71	8.0	13	1:1.9
	2006	7	4	0	57	10.7	9	1:1.3
	2007	7	1	0	14	30.0	18	1:2.6
	2008	7	3	0	43	4.5	6	1:1.0
	2009	4	0	0	0	ND	2	1:1.0
	2010	4	0	0	0	ND	4	1:1.0
	2011	4	1	0	25	40.0	1	1:1.0
17	2012	4	1	0	25	1.8	1	1:1.0
	2000 ^b	35	12	0	34	5.8	23	1:1.0
	2001	22	2	0	9	4.5	25	1:1.1
	2002	22	9	0	41	6.5	14	1:1.0
	2003	22	6	0	27	7.7	16	1:1.0
	2004	22	7	0	32	10.3	16	1:1.0
	2005	18	5	0	28	3.8	22	1:1.2
	2006	18	6	0	33	6.5	13	1:1.0
	2007	18	0	0	0	ND	18	1:1.1
	2008	18	5	0	28	8.5	17	1:1.0
	2009	5	1	0	20	15.0	7	1:1.4
	2010	5	1	0	20	1.0	2	1:1.0
	2011	5	1	0	20	1.0	1	1:1.0
	2012	5	0	0	0	ND	1	1:1.0
19	2000	14	7	0	50	5.6	29	1:2.1
	2001	12	2	0	17	14.0	15	1:1.3

Table 2. Continued.

Hunt area	Year	Tags	Harvest		Hunter success (%)	Days/hunter ^a	First-choice applicants	Drawing odds
			M	F				
20	2002	12	4	0	33	5.0	6	1:1.0
	2003	12	6	0	50	10.7	14	1:1.2
	2004	12	3	0	25	12.5	40	1:3.3
	2005	12	1	0	8	5.0	18	1:1.5
	2006	12	8	0	66	4.9	19	1:1.6
	2007	12	0	0	0	ND	19	1:1.6
	2008	12	3	0	25	6.7	7	1:1.0
	2009	5	1	0	20	5.0	3	1:1.0
	2010	5	1	0	20	2.0	1	1:1.0
	2011	5	0	0	0	ND	7	1:1.4
	2012	5	0	0	0	ND	9	1:1.8
	2000	14	5	0	36	11.4	23	1:1.6
	2001	14	5	0	36	8.4	17	1:1.2
	2002	14	4	0	29	4.5	14	1:1.0
	2003	14	2	0	14	7.0	10	1:1.0
	2004	14	2	0	14	16.5	9	1:1.0
	2005	10	3	0	30	17.5	8	1:1.0
	2006	10	2	0	20	12.0	12	1:1.2
	2007	10	3	0	30	4.0	11	1:1.1
	2008	10	2	0	20	15.0	6	1:1.0
2009	5	0	0	0	ND	3	1:1.0	
2010	5	0	0	0	ND	2	1:1.0	
2011	5	1	0	0	14.0	1	1:1.0	
2012	3 ^d	1	0	33	1.0	1	1:1.0	

^a Data from successful hunters only.

^b Some tags not sold.

^c 11 Permits were used, but 4 hunters elected to take rain checks (fire-related) and did not hunt

^d 5 Permits issued but 2 hunters elected to take rain checks (fire-related) and did not hunt

Table 3. Known moose mortalities, excluding controlled hunts, Clearwater Region, 1979-present.

Year	Mortality agent					Total
	Native American harvest	Illegal kill	Road kill	Natural	Other	
1979	4	9	4	0	0	17
1980	4	19	3	0	0	26
1981	1	13	4	0	0	18
1982	11	21	0	0	0	32
1983	13	25	5	0	0	43
1984	10	19	4	0	0	33
1985	6	15	4	0	0	25
1986	18	14	7	0	0	39
1987	2	13	11	0	0	26
1988	0	0	0	0	0	0
1989	4	17	7	0	0	28
1990	13	11	1	0	0	25
1991	15	21	3	0	0	39
1992	10	33	5	6	4	58
1993	7	31	5	0	2	45
1994	2	13	2	1	5	23
1995	10	4	7	4	2	27
1996	4	9	4	3	6	26
1997	1	18	2	2	5	28
1998	6	3	3	0	5	17
1999	6	1	0	0	8	15
2000	5	10	0	5	0	20
2001	1	9	3	0	1	14
2002	2	13	4	0	2	21
2003	0	2	0	0	3	5
2004	0	7	2	2	1	12
2005	2	7	6	2	0	17
2006	0	2	0	2	1	5
2007	1	2	1	0	1	5
2008	0	1	3	0	1	5
2009	1	2	3	0	0	6
2010	0	2	2	1	0	5
2011	0	0	2	1	1	4
2012	0	0	0	3	1	4

MAGIC VALLEY REGION

GMUs 44, 48, 49, 56, 73, 73A

Controlled Hunt Areas 44, 48, 56

Abstract

Moose populations have declined the northern portion of the Magic Valley Region and increased in the southern portion. Because of this, for the 2011 and 2012 seasons, antlerless harvest was eliminated and antlered tags reduced in GMUs 44, 48 and 49. Hunts were authorized for the first time in GMUs 55 and 57. In 2012, 8 total tags were issued in hunt areas 44 and 55 and all 8 hunters were successful.

Management Direction

Follow statewide management direction; allow established populations to expand; transplant moose where feasible; and increase effort to record sightings and mortalities.

Background

Prior to 1990, transient moose were recorded throughout the Magic Valley Region, but there were no viable, resident populations. From 1986-2000, 31 moose were released in GMUs 43 and 44. Following these releases, moose numbers in the region continued to increase as a result of good reproduction and natural ingress. Presently, viable populations capable of sustaining limited harvest occur in GMUs 44, 48, 49, 55, 56, and 57 (Figure 7).

Population Surveys

Aerial population surveys for moose have not been conducted in the region. During the 1990s and 2000s, increasing anecdotal observations indicated a growing moose population along the South Fork Boise River in GMU 43, Willow Creek in GMU 44, Big Wood River in GMU 48, and in the Trail Creek drainage on the border of GMUs 48-49. However, over the past few years, observations and harvest rates have declined. Populations in the Sublett area (GMU 56) appear to be stable and observations are common. In recent years, most have continued to expand their distribution and have become increasingly common in GMUs 54, 55, and 57.

Harvest Characteristics

Hunting season length for antlered moose in the 2 hunt areas in Magic Valley Region was 86 days in 2012 (Appendix A). Three antlered tags were offered in Hunt Area 44. The boundary of Hunt Area 44 was changed prior to the 2005 hunting season to include portions of GMUs 44 and 48. The boundary was again changed prior to the 2011 hunting season to include a portion of GMU 44 and all of GMUs 48 and 49. As a result, Hunt Area 48 was eliminated in 2011. All 3 bulls harvested in Hunt Area 44 in 2012 were taken in Unit 49 (Table 1).

Prior to 2011, Hunt Area 56 included GMUs 56, 73, and 73A. However, in 2011, because of growing moose populations in GMUs 55 and 57, GMU 56 was combined with those two GMUs into a new Hunt Area 55. Five antlered tags were offered in Hunt Area 55. Five bulls were

harvested, with 2 taken in GMU 55 and 3 taken in GMU 56 (Table 1). In addition, 2 Super Hunt tags holders harvest antlered moose in GMU 55.

No antlerless hunts were offered in the Magic Valley region during 2012.

Capture and Translocation

No moose were released in the region during this reporting period.

Management Implications

Moose population declines in the Big Wood River drainage and surrounding areas have resulted in the elimination of antlerless hunts and a reduction in antlered tags in Hunt Area 44 for the 2011 and 2012 seasons. Although no formal population surveys are planned, the population will continue to be monitored through incidental observations by agency personnel and the public.

Increasing moose numbers in the southern portion of the region (GMUs 54, 55, and 57) have provided the opportunity to expand hunting opportunities. GMUs 55 and 57 were opened to moose hunting for the first time in 2011 and were grouped with GMU 56 into Hunt Area 55. Consideration will be given in 2015 to opening Unit 54 to moose hunting as well.




Figure 7. Magic Valley and Southeast Antlered Moose Hunt Areas.

Table 1. Moose harvest and drawing odds by hunt area, Magic Valley Region, 1999-present.

Hunt area	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing odds
			M	F				
44 ^a	2001	2	2	0	100	3.8	9	1:4.5
	2002	2	1	0	50	1.0	13	1:6.5
	2003	4	3	0	75	11.0	16	1:4.0
	2004	4	4	0	100	7.7	20	1:5.0
	2005	6	2	0	33	6.5	13	1:2.2
	2006	6	1	2	50	6.5	21	1:3.5
	2007	6	3	1	67	3.5	10	1:1.7
	2008	6	1	1	33	5	23	1:3.7
	2009	6	1	1	33	19.5	18	1:3.0
	2010	6	1	0	17	5	11	1:1.8
	2011	3	2	0	67	4.5	17	1:5.6
	2012	3	3	0	100		11	1:3.7
48 ^b	2005	4	2	2	100	6.3	8	1:2.0
	2006	4	1	2	75	4.5	9	1:2.3
	2007	4	0	0	0	0	6	1:1.5
	2008	4	2	0	50	12.0	8	1:2
	2009	4	2	2	100	4.5	11	1:2.8
	2010	4	2	0	50	8.0	9	1:2.3
56 ^c	1999	5	5	0	100	16.0	28	1:5.6
	2000	5	5	0	100	3.8	21	1:4.2
	2001	5	4	1	100	19.2	31	1:6.2
	2002	5	4	0	80	3.0	31	1:6.2
	2003	5	5	0	100	17.2	37	1:7.4
	2004	5	5	0	100	5.6	44	1:8.8
	2005	5	5	0	100	12.3	46	1:9.2
	2006	5	5	0	100	4.5	42	1:8.4
	2007	5	5	0	100	7.8	73	1:14.5
	2008	5	3	0	60	10.0	114	1:22.8
55	2009	5	5	0	100		116	1:23.2
	2010	5	5	5	100	6.8	111	1:22.2
55	2011	5	5	0	100	7	138	1:27.6
	2012	5	5	0	100	2.4	97	1:19.4

^a Hunt established in 2001; includes portions of GMUs 44 and 48.

^b Hunt established in 2005; includes all of GMU 49 and a portion of GMU 48.

^c Prior to 2011 hunt area included GMUs 56, 73, and 73A. In 2011 and 2012 Hunt Area 55 was established and included GMUs 55, 56 and 57.

SOUTHEAST REGION

GMUs 66A, 70, 71, 72, 73, 73A, 74, 75, 76, 77, 78

**Controlled Hunt Areas 66A, 70, 71-1, 71-2, 72,
74, 75, 76-1, 76-2, 76-3, 77, 78**

Abstract

Eighty-two antlered-only and 45 antlerless-only tags were offered in 2012. Mandatory harvest reports identified a total of 63 antlered (76.8 % hunter success) and 27 antlerless (60.0 % hunter success) moose harvested. The average outside antler spread was 32.8 inches for antlered moose for which data were available.

Management Direction

Management direction for moose in Southeast Region follows that for the state in general: to provide “high-quality” hunting and other moose-related recreational opportunities. Consequently, tag levels are conservative, and hunter success is high relative to hunts for other cervid species. For antlered-only hunts, emphasis is on providing each hunter with the opportunity to harvest a mature bull moose. Antlerless-only moose hunting is also offered due to relatively high moose populations. Non-consumptive values of moose are also important.

The 1991-1995 Moose Management Plan established the goals of providing high-quality moose hunting and other moose-related recreational experiences for as many people as possible, assisting the expansion of moose populations into available habitat, and increasing tag numbers where possible.

Background

Prior to the 1950s, there were too few moose in Southeast Region to justify a harvest. The first hunt for moose in the region was held in 1959 when 5 antlered-only tags were issued for a portion of GMU 76. As moose populations continued to grow and expand to other areas in the Southeast Region so did the hunting opportunity and harvest, reaching a high of 180 moose harvested in 11 GMUs in 2003. Tag levels and total harvest have decreased significantly since 2003 in response to apparent changes in population densities. Illegal moose harvest may be substantial (Kuck and Ackerman 1984), although reporting of these cases is sporadic. The Department issued a small number of tags for any moose in several GMUs from 1975-1990. An average of 80% of that harvest was antlered moose. In 1991, antlerless-only hunts were instituted in GMUs 66A and 76. Since 1991, tags have been issued for antlered or antlerless-only moose (Figures 8 & 9). Antlerless moose hunts start later than antlered hunts to provide more time for calf development.

Population Surveys

No moose surveys were conducted in the Southeast Region during the reporting period. However, moose were counted incidentally during an elk survey in GMU's 76 and 66A in late winter 2013. A total of 122 moose were observed (52 bulls, 43 cows, 23 calves, 4 unknown) with a calf:cow:bull ratio of 54:100:121. Because this was not a randomized aerial sightability

survey designed specific for moose caution must be taken interpreting the results as there is bias in the data; however, this information does provide some general baseline trend information for future comparison.

The most recent moose specific survey was conducted January 2002 in GMUs 66A and 76-3. In Hunt Area 66A, 19 search units were stratified as high, medium, or low likelihood of moose and 13 search units were flown for sightability. One hundred fifty-two moose were counted in these 13 search units consisting of 75 cows, 48 bulls, and 29 calves (Table 1). Estimates of 219 (± 31) total moose including 105 (± 15) cows, 75 (± 18) bulls, and 39 (± 9) calves were generated using the Hiller-Soloy Wyoming-based model (Unsworth et al. 1994). Overall herd composition was estimated as 48% cows, 34% bulls, and 18% calves. The population estimate of 219 in 2002 was 23% lower than the estimate of 285 in 1995; however, 90% confidence intervals overlap. The average number of moose observed was 3.0 in low units, 16.0 in medium units, and 18.5 in high units. Search units were likely well-stratified for the survey.

During the 2002 survey, in Hunt Area 76-3, 13 search units were stratified as high or low likelihood of moose and 10 search units were flown for sightability. One hundred three moose were counted in these 10 search units consisting of 41 cows, 48 bulls, and 14 calves (Table 1). Estimates of 174 (± 40) total moose including 71 (± 20) cows, 78 (± 20) bulls, and 25 (± 8) calves were generated using the Hiller-Soloy Wyoming-based model. Overall herd composition was estimated as 41% cows, 45% bulls, and 14% calves. The population estimate of 174 in 2002 was very close to the 167 estimated in 1995. The average number of moose observed was 9.8 in low units and 11.2 in high GMUs. Search areas may need to be re-stratified or have stratification by moose likelihood deleted in future surveys.

Harvest Characteristics

Tag levels (Tables 2 and 3) for 2012 remained the same as those in 2011. However, current tag levels are reduced from what was available in 2010 due to concerns over declining population numbers in some areas. The 160 tags (95 antlered and 65 antlerless) in 2010 were reduced to 127 tags (82 antlered, 45 antlerless) in 2011. Minimum reported harvest was available through a mandatory mortality report of successful hunters. Reported harvest totaled 90; 63 antlered and 27 antlerless moose (Tables 2 and 3). Average antler spread for Southeast Region was 32.8 inches.

Minimum overall hunter success rate for the region was 69.2 %; 60.0 % for antlerless-only tags and 76.8 % for antlered-only tags.

Other sources of moose mortality are Native American harvest, natural, road-kills, illegal, and other. For the 2010-2011 reporting period, 15 non-harvest mortalities were reported (Table 4). Reporting of non-hunting mortalities is believed to be much lower than the actual number.

Climatic Conditions

Winter 2011-2012 snow depths averaged 65 – 68 % of the 30-year mean for most of the southeast region. Spring conditions were mild with below average moisture levels extending into the summer months.

Habitat Conditions

Succession of aspen stands into conifer may negatively affect moose habitat in the future. Treatment to retard succession may slow potential decreases. Development and disturbance associated with mining and timber harvest in the eastern portion of the region continues to threaten moose habitats. Livestock grazing and other development of riparian areas impacts moose habitat in many parts of the region.

Management Implications

The mandatory check of moose harvested provides the majority of information available for management. Aerial surveys, using sightability models such as Anderson (1994) and Unsworth et al. (1994), provided solid background data in 2002 and years prior, but limitations on resources has not allowed for repetition of these surveys since. Currently, the only population data obtained on moose is from incidental observations during mule deer and elk aerial surveys. As such, tag levels are set at conservative levels that likely allow for passive population expansion and growth, particularly in those areas being newly colonized.

The drawing odds for antlered-only tags indicate strong demand for moose hunting opportunity. Drawing odds varied considerably in 2012, with hunts being as high as 1:17.8 and as low as 1:2.4, depending on the hunt. Antlerless-only drawing odds are generally 1:1 or less; however, leftover tags sell quickly.

Moose also have high non-consumptive values for viewing by the public. Their relative abundance and general lack of fear of humans when in the open make them easy for people to observe.

Moose translocations and hazing activities are expanding to include the entire year rather than spring and early summer. Over the course of a year, on average, 5 – 30 moose wander into the city of Pocatello and surrounding communities. These are nearly always yearlings or 2-year olds and are most often hazed back into the surrounding hills or captured using chemical immobilization and translocated to more suitable habitat.

Moose population data may need to be collected again in the form of specific sightability surveys or incidentally during deer and elk surveys in the future. Wyoming is experiencing unexplained declines in moose populations directly to the east of the Southeast Region. Some possible explanations may be parasites such as carotid artery worm (which has been documented in Idaho moose and in this region), disease, climate change, or alterations in moose habitat.

Literature Cited

Anderson, C. R. 1994. Aerial moose sightability in western Wyoming. Thesis, University of Wyoming, Laramie, USA.

Kuck, L., and B. Ackerman. 1984. Impacts of illegal harvest on big game. Pages 363-373 in L. Kuck (ed.) Cooperative Wildlife Study, Phase 2: Mining Impacts Studies. Idaho Department of Fish and Game, Boise, USA.

Unsworth, J. W., F. A. Leban, D. J. Leptich, E. O. Garton, and P. Zager. 1994. Aerial survey: user's manual. Second edition. Idaho Department of Fish and Game, Boise, USA.




Figure 8. Southeast Antlered Moose Hunt Areas.

Table 1. Total observed moose by sex/age class and model estimates of moose from aerial surveys, Southeast Region, 1991-2002.

Hunt area Year	Observed		Estimate	
	Total	Bull:cow:calf	Total	Bull:cow:calf
76-1, 2				
1994	90	42:100:42	432	26:100:50
2000	286	74:100:42	510±83	74:100:42
76-3, 4				
1993	104	76:100:37	192	76:100:36
1997	89	85:100:44	190	100:100:53
76-5, 6				
1991	136	49:100:60		
1995	121	55:100:40	167±22	54:100:34
2002	103	117:100:34	174±40	110:100:35
76				
1999	140	100:100:62	583±146	99:100:60
66A				
1995	159	69:100:49	285±60	67:100:43
2002	152	64:100:39	219±31	71:100:37

Table 2. Moose harvest and drawing odds, Southeast Region, 1984-present.

Year	Tags	Harvest			Hunter success (%)	First-choice applicants	Drawing odds
		M	F	Total			
1984	95	77	5	82	86	1,908	1:20.1
1985	95	73	4	77	81	1,841	1:19.4
1986	95	79	4	83	87		
1987	95	81	8	89	94	834	1:8.8
1988	110	100	5	105	95	830	1:7.5
1989	110	95	4	99	90	556	1:5.1
1990	125	98	9	107	86	738	1:5.9
1991	135	94	20	114	84	910	1:6.7
1992	135	98	19	117	87	837	1:6.2
1993	160	113	29	142	89	728	1:4.6
1994	160	114	29	143	89	809	1:5.1
1995	180	115	32	147	82	932	1:5.2
1996	180	105	34	139	77	921	1:5.1
1997	180	115	31	146	81	849	1:4.7
1998	180	103	28	131	73	804	1:4.5
1999	185	104	49	153	83	1,026	1:5.5
2000	185	111	34	145	78	600	1:3.2
2001	220	124	48	172	78	747	1:3.4
2002	220	127	38	165	75	723	1:3.3
2003	225	129	51	180	80	701	1:3.1
2004	225	129	31	160	71	737	1:3.1
2005	160	75	41	116	73	736	1:4.6
2006	160	81	40	121	76	647	1:4.0
2007	160	80	39	119	74	715	1:4.5
2008	160	72	37	109	68	667	1:4.2
2009	160	80	44	124	78	809	1:5.0
2010	160	71	36	107	67	696	1:4.4
2011	127	63	23	86	68	788	1:6.2
2012	127	63	27	90	71	623	1:4.9

Table 3. Moose harvest and drawing odds by hunt area, Southeast Region, 2000-present.

Hunt area	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing odds
			M	F				
66A	2000	42	27	7	81	5.7	194	1:4.6
	2001	45	24	12	80	4.1	220	1:4.9
	2002	45	29	12	91		202	1:4.5
	2003	45	28	12	89	3.8	215	1:4.8
	2004	45	30	7	82	6.5	197	1:4.8
	2005	25	15	8	92	4.1	188	1:7.5
	2006	25	14	9	92	4.5	176	1:7.0
	2007	25	10	6	64	7.2	170	1:6.8
	2008	25	12	8	80	4.7	131	1:5.2
	2009	25	13	8	84	4.6	151	1:6.0
	2010	25	14	5	76		147	1:5.9
	2011	15	9	3	80		129	1:8.6
2012	15	9	3	80		85	1:5.7	
70	2000	5	4	0	80	20.0	21	1:4.2
	2001	5	4	0	80	11.8	15	1:3.0
	2002	5	5	0	100		30	1:6.0
	2003	5	5	0	100	10.0	15	1:3.0
	2004	5	5	0	100	5.8	34	1:3.0
	2005	5	4	0	80	10.0	47	1:9.4
	2006	5	5	0	100	3.6	68	1:13.6
	2007	5	5	0	100	10.5	75	1:15.0
	2008	5	5	0	100	10.8	50	1:10.0
	2009	5	4	0	80	4.0	99	1:19.8
	2010	5	5	0	100		68	1:13.6
	2011	5	4	0	80		105	1:21.0
2012	5	5	0	100		89	1:17.8	
71	2000	15	7	4	73	11.0	42	1:2.8
	2001	20	9	5	70	7.1	54	1:2.7
	2002 ^a	20	7	3	50		25	1:1.3
	2003 ^a	20	9	6	75	7.5	23	1:1.2
	2004	20	8	3	55	4.1	34	1:1.2
	2005	20	6	3	45	8.0	34	1:1.2
	2006	20	8	6	70	8.2	36	1:1.8
	2007	20	8	7	75	2.5	45	1:2.3
	2008	20	6	4	50	7.0	52	1:2.6
	2009	20	6	7	65	5.8	58	1:2.9
	2010	20	2	6	40		25	1:1.3
	2011	20	7	4	55		32	1:1.6
2012	20	3	4	35		39	1:2.0	
72	2000	5	5	0	100	5.4	26	1:5.2
	2001	5	5	0	100	1.8	39	1:7.8
	2002	5	5	0	100		31	1:6.2
	2003	5	4	0	80	12.8	34	1:6.8

Table 3. Continued.

Hunt area	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing odds
			M	F				
74	2004	5	5	0	100	6.8	27	1:6.8
	2005	5	5	0	100	5.6	27	1:6.8
	2006	5	5	0	100	15.6	33	1:6.6
	2007	5	4	0	80	11.8	34	1:6.6
	2008	5	5	0	100	12.2	41	1:8.2
	2009	5	5	0	100	10.6	36	1:7.2
	2010	5	5	0	100		44	1:8.8
	2011	5	4	0	80		35	1:7.0
	2012	5	2	0	40		35	1:7.0
	2000	5	4	0	80	13.7	12	1:2.4
	2001	5	4	0	80	34.7	16	1:3.2
	2002	5	3	0	60		16	1:3.2
	2003	5	4	0	80	7.0	24	1:4.8
	2004	5	3	0	60	13.7	17	1:4.8
	2005	5	5	0	100	6.0	22	1:4.4
	2006	5	4	0	80	10.5	21	1:5.3
	2007	5	5	0	100	10.4	23	1:4.6
	2008	5	3	0	60	12.0	22	1:4.4
	2009	5	5	0	100	16.5	29	1:5.8
	2010	5	5	0	100		34	1:6.8
2011	5	4	0	80		33	1:6.6	
2012	5	4	0	80		21	1:4.2	
75	2000	15	5	4	60	3.8	28	1:1.9
	2001	15	10	4	93	7.1	26	1:1.7
	2002	15	9	2	73		29	1:1.9
	2003 ^a	15	9	3	80	6.8	31	1:2.1
	2004	15	9	3	80	8.1	36	1:2.1
	2005	10	3	3	60	10.0	30	1:3.0
	2006	10	4	4	80	5.4	42	1:4.2
	2007	10	5	3	80	3.6	26	1:2.6
	2008	10	4	4	80	11.4	40	1:4.0
	2009	10	4	4	80	6.9	71	1:7.1
	2010	10	4	3	70		51	1:5.1
	2011	10	4	4	80		47	1:4.7
2012	10	4	1	50		49	1:4.9	
76	2000	84	45	19	76	5.6	249	1:3.0
	2001	105	51	27	74	4.8	326	1:3.1
	2002 ^a	105	57	21	74		329	1:3.1
	2003	110	51	30	74	6.2	323	1:2.9
	2004	110	51	18	63	6.9	321	1:2.9
	2005	70	28	20	69	4.8	335	1:4.8
	2006	70	28	14	60	6.3	211	1:3.0
	2007	70	32	15	78	6.7	290	1:4.1

Table 3. Continued.

Hunt area	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing odds
			M	F				
77	2008	70	28	13	59	6.7	270	1:3.9
	2009	70	36	14	71	6.9	299	1:4.3
	2010	70	24	16	57		231	1:3.3
	2011	42	18	8	62		210	1:5.0
	2012	42	22	12	81		143	1:3.4
	2000	7	7	0	100	7.1	12	1:1.7
	2001	10	8	0	80	7.6	24	1:2.4
	2002	10	4	0	40		25	1:2.5
	2003	10	9	0	90	6.3	23	1:2.3
	2004	10	9	0	90	5.4	20	1:2.3
	2005	10	5	3	80	11.4	23	1:2.3
	2006	10	5	5	100	6.1	34	1:3.4
	2007	10	5	3	80	6.7	28	1:2.8
	2008	10	4	4	80	15.1	38	1:3.8
78	2009	10	5	3	80	8.5	29	1:2.9
	2010	10	4	3	70		41	1:4.1
	2011	10	2	1	30		37	1:3.7
	2012	10	4	3	70		25	1:2.5
	2000	7	7	0	100	13.9	16	1:2.3
	2001	10	9	0	90	10.9	27	1:2.7
	2002	10	8	0	80		36	1:3.6
	2003	10	9	0	90	19.8	13	1:1.3
	2004	10	9	0	90	8.2	51	1:1.3
	2005	10	4	4	80	20.3	30	1:3.0
	2006	10	5	2	70	4.4	26	1:2.6
	2007	10	5	4	90	5.5	24	1:2.4
	2008	10	5	4	90	5.4	23	1:2.3
	2009	10	4	5	90	7.4	37	1:3.7
2010	10	5	5	100		45	1:4.5	
2011	10	5	4	90		64	1:6.4	
2012	10	5	5	100		51	1:5.1	

^a Applicants and drawing odds for antlered hunts only.

Table 4. Known moose mortalities, excluding controlled hunts, Southeast Region, 1993-present.

Year	Mortality agent						Total
	Native American harvest	Illegal kill	Road kill	Natural	Train kill	Other	
1993	0	0	2	0	0	0	2
1994	0	0	1	0	0	0	1
1995	1	10	1	1	0	7	20
1996	1	2	5	0	1	1	10
1997	0	1	1	3	0	3	8
1998	0	1	1	0	1	3	6
1999	0	1	4	3	0	0	8
2000	0	4	2	1	0	2	9
2001	1	1	3	0	0	4	9
2002	0	1	2	1	0	1	5
2003	0	0	2	3	0	1	6
2004	0	0	2	1	0	0	3
2005	0	1	2	0	0	0	3
2006	0	0	1	3	0	0	4
2007	0	1	1	5	0	0	7
2008	0	1	1	3	0	1	6
2009	0	0	4	1	0	0	5
2010	0	0	0	0	0	1	1
2011	0	2	3	1	0	3	9
2012	1	0	7	2	0	5	15

UPPER SNAKE REGION

Abstract

Hunting season lengths for antlered and antlerless moose remained at 86 days (30 Aug-23 Nov) and 40 days (15 Oct-23 Nov), respectively, in 2012 (Figures 10 & 11). Due to concerns over hunter success and/or bull quality, tag numbers for the 2009 season were reduced significantly from the 2008 levels. Twenty one controlled hunts with 182 tags were offered for antlered moose and 10 controlled hunts with 55 tags were offered for antlerless moose in the Upper Snake Region in 2012 (Table 1). A total of 156 antlered (80% hunter success) and 45 antlerless (82% success) moose were harvested in 2012 as determined by BGMR reports. The mean antler spread for all antlered hunts combined was 37.5 inches. Overall drawing odds for moose hunts were 1:5.9 (Table 1).

Other sources of moose mortality are Native American harvest, natural, road-kill, train-kill, illegal, and other. Ten non-harvest mortalities were reported for the Upper Snake Region in 2012 (Table 2).

No population surveys were conducted specifically for moose during this reporting period due to fiscal constraints.

Climatic Conditions

Overall, climatic conditions were average for moose throughout this reporting period. The summer of 2012 was drier than average. The winter of 2012-2013 was mild, with less than average snow pack and crusting.

Depredation, Capture, and Translocation

Nuisance moose complaints in and around houses and towns are common in the Upper Snake Region and are often dealt with through hazing, public education, or relocation of the animal. Due to the mild winter conditions in the region during 2012-2013, there were few nuisance moose complaints. However, some moose have to be moved from human habitation due to conflicts and human safety concerns. Regional staff chemically immobilized 2 moose and relocated them to suitable, occupied moose habitat within the Upper Snake Region this reporting period.




Figure 10. Upper Snake and Salmon Antlered Moose Hunt Areas.




Figure 11. Southeast and Upper Snake Antlerless Moose Hunt Areas.

GMUs 50, 51, 58, 63, 63A
Controlled Hunt Areas 50, 51, 63, 63A

Background

In early 1980, 6 moose were released near North Fork of the Big Lost River (GMU 50). Most initially remained close to their release site, but there has been egress to other areas. Reproduction occurred, and additional transplants augmented this population. During winter 2001-2002, several nuisance moose were also translocated to GMU 50.

An antlered hunt in GMU 50 was initiated in 1993 and an antlerless hunt was initiated in 2003. An antlered moose hunt was opened in GMU 51 in 1999 as a result of an increasing number of moose being sighted incidentally during deer and elk sightability surveys and ground observations. In 2003 and 2004, an antlered hunt was authorized in GMU 58 for the same reason but was subsequently closed in 2005.

A significant population of moose exists in GMU 63A. Moose utilize riparian habitat along the North and South Forks of the Snake River and associated sloughs, and depredation and nuisance complaints occur on a fairly regular basis. Moose distribution in GMU 63 is centered around the Mud Lake Wildlife Management Area (WMA)-Camas National Wildlife Refuge (NWR) area.

Hunt Area 63A was initiated in 1987. GMU 63 was added to Hunt Area 63A in 1999 and was then split into 2 separate hunts (Hunt Areas 63 and 63A) in 2003. Due to declines in harvest success, average antler spread, and moose observed during Mud Lake WMA spotlight surveys, all moose harvest was eliminated in Hunt Area 63 for the 2009 and 2010 seasons.

Population Surveys

No moose population surveys were conducted during this reporting period.

Harvest Characteristics

A total of 42 tags were issued in these GMUs in 2010, resulting in the harvest of 32 animals (80% success) based on mandatory harvest reports (Table 3). Mean antler spreads were 38 ($n = 5$) in GMU 50, 36 ($n = 3$) in GMU 51, and 40 ($n = 13$) in GMU 63A.

Habitat Conditions

Habitats within these GMUs are quite varied. In GMU 50, extensive willow bottoms provide good summer and winter habitat, and the moose population appears to be increasing and ranging throughout the coniferous zone in summer. Habitat in GMUs 51 and 58 are limited to discontinuous willow riparian areas. Habitat in GMU 63 is almost entirely desert and unsuitable for moose, except areas on and adjacent to Mud Lake WMA and Camas NWR. Habitat in GMU 63A consists primarily of the Snake River riparian zone adjacent to private residential and agricultural lands.

Management Implications

Based on harvest data, populations in most hunt areas appear to be relatively stable. However, there may need to be changes made in Hunt Area 51 moose opportunity to address declines in harvest success and average antler spread. The opportunity to reinstate moose harvest in 63A will be re-evaluated during the 2013-2014 moose season setting process.

GMUs 59, 59A

Controlled Hunt Area 59

Background

Former Hunt Areas 59 and 59A were combined in 1993 to form the current Hunt Area 59. Prior to 1993, 2 hunts with a total of 12 antlered tags were offered in these GMUs. Former Hunt Area 59 had been open continuously since 1974 with tag levels fluctuating between 4 and 8 with over 90% hunter success reported. Hunt Area 59A was closed in 1978 after only 1 moose was harvested in the preceding 4 years. In 1983, this hunt was reopened and 2 tags were issued annually through 1988 with 100% hunter success. Four tags were issued each season from 1989-1992 with 100% hunter success. Due to declines in average antler spread and harvest success, combined with concerns from sportsmen and field personnel, harvest opportunity was significantly reduced in Hunt Area 59 for the 2009 and 2010 seasons. During the 2005-2008 seasons, there were 15 antlered and 5 antlerless tags available in this hunt. For the 2009 and 2010 seasons, tags were reduced to 5 antlered and no antlerless.

Population Surveys

A moose trend count was flown in GMUs 59 and 59A on 17-18 December 1994 using a Bell Model G47 Soloy helicopter. Counting conditions were good, with 8 or more inches of relatively new snow cover present over the entire area. All probable moose habitat was surveyed. A total of 179 moose (129 in GMU 59 and 50 in GMU 59A) with a bull:cow:calf ratio of 44:100:54 was counted on the survey. Of the 40 bulls counted, 13 were classified as yearlings, 20 as adults, and 7 had already shed antlers.

Few previous data are available for comparison. Prior to this count, no surveys had been conducted in GMU 59 since 1984 (64 total moose), and GMU 59A had never been surveyed specifically for moose. However, during deer and elk sightability surveys, moose were counted on an incidental basis. In 1991-1992, 46 moose were counted in GMU 59 and 71 in GMU 59A. In 1993-1994, 49 moose were observed in GMU 59 and 46 in GMU 59A (unclassified). The 1999-2000 survey resulted in a total count of 90 moose (10 bulls, 19 cows, 13 calves, 48 unclassified). The 2004-2005 survey resulted in a total count of 74 moose (6 bulls, 13 cows, 6 calves, 49 unclassified). During the 2008 Beaverhead elk survey, 11 and 13 moose were incidentally counted in GMUs 59 and 59A, respectively. Not all moose habitat is flown during elk surveys, so these incidental numbers are not a reliable estimate of the number of moose in an area. No aerial surveys were conducted in this Hunt Area during the reporting period.

Harvest Characteristics

Five tags for antlered moose were offered in 2012, hunter success was 100% (Table 3) and mean antler spread was 33 inches ($n = 5$).

Known illegal kill (Table 2) was a serious problem in the early 1980s when it nearly equaled controlled harvest but has been of little significance, based on documented mortalities, in recent years.

Habitat Conditions

Habitat consists primarily of conifer/sagebrush ecotones and aspen. Riparian areas are limited and discontinuous. Habitat extends down major drainages that have willows. Improving riparian zone management would increase habitat quality and quantity in this area.

Management Implications

General observations indicate the moose population in these GMUs has declined in recent history. Additionally, average antler spread on harvested bulls was below the management objective of 35 inches during the 2005-2008 seasons. Therefore, tags were significantly reduced for 2009 and 2010 in an attempt to increase total numbers and bull quality in the Hunt Area. The effects of the tag reduction on average antler spread and harvest success should continue to be monitored.

GMUs 60, 60A, 61, 62, 62A

Controlled Hunt Areas 60, 60A, 61-1, 61-2, 61-3, 62, 62A

Background

During the 1970s, the moose population in Fremont County was thought to be declining and experiencing high levels of illegal mortality and Native American harvest. As a result, all moose hunts in Fremont County were closed in 1977. After a boundary change to include only Clark County, Hunt 61-1 was the only hunt open from 1977 to 1982.

A winter aerial survey conducted in 1983 counted moose in numbers slightly below the highs of the early 1950s. The Island Park area is the only area where counts were clearly lower than those in the 1952-1956 periods. In response to the population recovery, 8 controlled hunts were opened in 1983 in Fremont County.

A new hunt was established in GMU 60A in 1986. The hunt area consists of agricultural land and the riparian zone along Henrys Fork of the Snake River. Many residences and farms are in the area. The moose population within this corridor has been increasing. We received many depredation and nuisance complaints of moose in agriculture fields and near towns and residences, resulting in expanded antlerless hunting opportunity. Tags were reduced by approximately 50% on the Island Park caldera portion of the region in 1991 as a result of significant winter mortality during the winter of 1988-1989, but steadily increased through 2004

as the populations continued to grow. Like other portions of the region, tag levels were significantly reduced during 2005-2007 in an attempt to increase the number of larger bulls in the population.

Based on declines in harvest success and average antler spread in many of these Hunt Areas, along with concerns over the unknown effect of an expanding wolf population on moose numbers, antlered and antlerless tags were reduced within these hunts for the 2011 and 1012 seasons. A total of 55 antlered and 10 antlerless tags were offered in 2011, which was a 15% and 33% reduction in antlered and antlerless tags, respectively.

Population Surveys

A population survey was conducted in GMU 62 and a portion of 62A during December 2000. The survey in 62A was not completed because of fiscal constraints. The final population estimate for GMU 62 was 366 moose including 180 cows, 109 bulls, and 77 calves (Table 4). This total compares to fixed wing censuses of 228 cows and 97 bull moose observed during 1989 and 1990, respectively.

Most of the area was surveyed by airplane from November 1989-February 1990 (Table 5). Survey results indicated that moose populations had decreased substantially since the previous winter. Moose appeared to be in poor condition prior to the 1988-1989 winter, following 2 years of drought, and significant winter losses probably occurred. In 2011, 125 moose were counted incidental to the Teton elk survey (portions of GMUs 65, 62, and 62A).

A helicopter survey was conducted along the North Fork Snake River corridor between St. Anthony and the Highway 33 Bridge in Hunt Area 60A in December 1991. Only the riparian corridor was searched, so this should be considered a minimum count. A total of 37 moose were observed (2 bulls, 21 cows, 14 calves).

Moose have been counted incidental to deer and elk sightability surveys in GMU 60A on a fairly regular basis. However, moose distribution varies greatly from year to year and, since not all search units are surveyed, the usefulness of this information is questionable.

In 2010, a total of 241 moose (104 cows, 61 bulls, 55 calves, 21 unclassified) were counted incidental to an elk sightability survey in GMU 60A (and small portions of GMUs 60, 61, and 62A). This was the first time in recent history an effort was made to document composition of the moose seen incidentally to an elk or deer survey. Other recent totals for GMU 60A (most unclassified) include 328 (2007), 239 (2004), 185 (2003), 387 (2002), 473 (2000), 585 (1998), 340 (1997), 219 (1996), 272 (1996), 360 (1995), 187 (1994), and 312 (1993). Twenty-two moose were counted incidental to the 2010 elk survey in GMUs 62 and 65 east (12 cows, 6 bulls, 4 calves) and 38 moose (unclassified) were counted during 2007 deer trend surveys in GMU 62.

Harvest Characteristics

Sixty-five moose tags were issued in 2012, resulting in the harvest of 58 animals (89% success) based on BGMR reports (Table 3). Mean antler spreads were 40 ($n = 13$) in GMU 60, 39 ($n = 4$) in GMU 60A, 36 ($n = 23$) in GMU 61, 35 ($n = 4$) in GMU 62, and 39 ($n = 4$) in GMU 62A.

Management Implications

The increase in desert-wintering moose has led to increased depredations and nuisance complaints during average to severe winters. Mortality during the 1988-1989 winter resulted in significant population declines. However, moose populations have rebounded rapidly to levels above those present prior to the 1988-1989 die-off. Consequently, tag levels were increased accordingly through 2004. Populations appear to be relatively stable, but mean antler spread and harvest success declined in many Hunt Areas between 2007 and 2009. The influence of wolves on the moose population in the Island Park caldera is not well understood. Therefore, harvest metrics should continue to be monitored carefully into the future. Research designed to investigate the relationship between wolves and moose in this area would be beneficial to effective moose management.

GMUs 64, 65, 67

Controlled Hunt Areas 64, 65, 67-1, 67-2

Background

All of GMU 64 except the Canyon Creek drainage, GMU 65, and GMU 67 north and west of State Highway 31 has been open to moose hunting since 1974. In 1983, this area (old Hunt Area 364) was split along GMU boundaries into 3 separate hunts. Increasing moose populations allowed a steady increase in tag levels until 1987. A new Hunt Area, 67-2, was created in 1983 and allowed the harvest of moose in that portion of GMU 67 previously closed. An antlerless hunt with 5 tags was created in 2005 in GMU 65.

Hunting opportunity has increased in these GMUs from 1 hunt with 2 tags during the early 1980s to 7 hunts with 78 tags (58 antlered and 20 antlerless tags) in 2004. Tags were subsequently reduced in 2005 to 65 (45 antlered and 20 antlerless) and have remained at this level since.

Population Surveys

Historically, moose populations appeared to be increasing in these GMUs prior to the winter of 1988-1989. Forage was impacted by 2 years of drought and moose shifted their distribution to lower elevation agricultural and urban areas. Moose appeared to be in poor condition and significant winter losses likely occurred.

During winter 1992-1993, moose were first counted incidental to elk sightability surveys. Totals of 48, 26, and 90 moose were counted in GMUs 64, the western portion of 65, and 67, respectively. Most animals counted were unclassified. Moose were also counted incidental to elk sightability surveys during the 1995-1996 winter. Totals of 36, 101, and 60 moose were observed in GMUs 64, 65, and 67, respectively. Again, most animals were not classified. Moose were again counted incidentally during the 1997-1998 winter. Totals of 67, 30, and 88 (largely unclassified) moose were counted in GMUs 64, western 65, and 67, respectively. Moose were counted in GMUs 64, 65, and 67, incidental to elk surveys during the 2003-2004 winter and a total of 110 moose were observed. In 2007, a total of 38 moose were counted in

GMU 64 incidental to mule deer trend surveys. During 2008, 139 moose were counted incidental to the Palisades elk survey (31 in GMU 64, 43 in GMU 65, and 65 in GMU 67). A total of 237 (59 in GMU 64, 107 in GMU 65, and 71 in GMU 67) moose (unclassified) were counted incidental to the Palisades deer survey in 2010.

Harvest Characteristics

Hunters harvested 49 moose on 61 tags (80% hunter success) in 2012 (Table 3). Mean antler spreads were 38 ($n = 10$) in GMU 64, 39 ($n = 9$) in GMU 65, and 41 ($n = 12$) in GMU 67.

Habitat Conditions

Conifer with interspersed aspen and narrow riparian areas make up the majority of moose habitat in this area. Mountain mahogany on south-facing ridges provides important winter moose habitat in GMUs 65 and 67. In GMU 64, moose are found wintering primarily in stream bottom willow/aspen/dogwood communities.

Management Implications

A 1989 aerial survey found approximately half the number of moose counted in 1985. A shift in moose distribution resulting from drought and severe winter conditions was partially responsible for the low count. Also, mortality during the 1988-1989 winter was above normal. Tag levels were maintained for the 1989 and 1990 seasons, but were adjusted in 1991 in response to data analysis. Moose populations appear to have rebounded rapidly to levels at or above those present prior to the 1988-1989 die-off. Consequently, tag levels increased in 1993, 1995, 1997, and again in 1999. Additionally, an antlerless hunt was initiated in GMU 64 in 1993. Bull tags were reduced, starting in 2005, in an attempt to increase the number of larger bulls in the population. In recent years, the moose population appears relatively stable (based on incidental counts) and the bulls harvested in GMUs 65 and 67 have had the highest average antler spread in the Upper Snake Region.

GMUs 66, 69

Controlled Hunt Areas 66-1, 66-2, 69-1, 69-2, 69-3

Background

The moose population in these GMUs increased at a fairly rapid rate during the late 1970s when populations elsewhere in Upper Snake Region were decreasing or remaining static. Hunts 66 and 69 were split in 1981 to create 4 hunts (66-1, 66-2, 69-1, and 69-2). This resulted in a 50% increase in tag levels from 1980 (16 to 24). A new hunt (69-3) was created in 1984 from adjacent portions of Hunts 66-1 and 69-2.

Hunt 69-1 was changed from antlered-only to either-sex in 1986 to address landowner concerns over grain field depredations. Either-sex tags were not effective in harvesting antlerless moose; no female moose were harvested. As a result, this hunt was changed back to antlered-only in 1991. However, beginning in 1993, an antlerless hunt (69-4) was initiated. This hunt had 10 tags and included all of GMU 69. In 1999, GMU 66 was added to this hunt, tags were increased

to 20, and it was renumbered Hunt Area 66-3. This antlerless hunt was restructured again in 2001. GMU 66 was dropped from the hunt area and GMU 69 was split into 3 hunt areas (69-1, 69-2, and 69-3) that correspond to the like-numbered antlered hunts.

Average antler spread of bull moose harvested in GMU 66 from 2004-2008 was below the management goal of 35 inches. Therefore, both antlered and antlerless tags were reduced (antlerless tags eliminated) for the 2009 and 2010 seasons to increase moose numbers and increase trophy quality of bulls. Seven hunts with 55 antlered tags and 10 antlerless tags were offered in GMUs 66 and 69 during 2011, compared with 60 antlered tags and 15 antlerless tags that were offered from 2009-2010. This was a 8% reduction in antlered and a 33% reduction in antlerless tag levels.

Population Surveys

No population surveys have been conducted in these GMUs specifically to monitor moose populations. However, moose were counted incidentally during deer and elk sightability surveys (not all subunits containing moose were surveyed). A total of 276 moose (131 cows, 89 bulls, 59 calves) were counted incidentally to the Tex Creek elk survey (GMUs 66 and 69) in 2010. Other recent totals, during various deer and elk surveys, include 169 (2008), 304 (2007), 384 (2005), 317 (2000), 228 (1999), 293 (1997), 200 (1995), 98 (1994), and 147 (1992).

Harvest Characteristics

Seven hunts with a total of 65 tags were offered in these GMUs in 2012 (Table 3). A total of 57 moose were harvested on 65 tags (88% success). Mean antler spreads were 34 ($n = 18$) in GMU 66 and 36 ($n = 33$) in GMU 69.

Habitat Conditions

GMU 66 is characterized by conifer/aspen habitats with narrow canyon bottom riparian areas which support moderate willow/dogwood communities. GMU 69 is primarily aspen/sagebrush and private agricultural land with willow riparian areas in most canyon bottoms. Tex Creek WMA contains important winter habitat for a variety of ungulates, including moose, and moose from adjacent areas may be migrating to the WMA to winter.

Management Implications

Steadily increasing moose populations in these GMUs resulted in an increase in tag levels in all of these hunts from the early 1990's through 2005. Additionally, an antlerless hunt has been offered since 1993. Bull tags were reduced, starting in 2005, in an attempt to increase the number of larger bulls in the population. Mean antler spread of bulls harvested remained slightly below the management goal of 35 inches from 2004-2008, in most Hunt Areas. GMUs 66 and 69 have the habitat conditions needed to produce trophy-class bulls. In an effort to increase bull quality, a number of bull and cow tags in these GMUs (particularly in GMU 66) were eliminated during the 2009-2010 trophy species season setting process. These changes should continue to be monitored to evaluate their effectiveness in increasing trophy bull quality.

Table 1. Moose harvest and drawing odds, Upper Snake Region, 1982-present.

Year	Tags	Harvest			Hunter success (%)	First-choice applicants	Drawing odds (%)
		M	F	Total			
1982	42	35	0	35	83	2,434	2
1983	88	86	0	86	98	3,357	3
1984	98	96	0	96	98	3,049	3
1985	120	118	0	118	98	3,403	4
1986	145	143	1	144	99	2,071	7
1987	148	144	2	146	99	1,970	8
1988	140	134	2	136	97	1,597	9
1989	145	129	6	135	93	1,248	12
1990	148	143	2	145	98	1,204	12
1991	128	111	14	125	98	1,554	8
1992	128	109	16	125	98	1,162	11
1993	214	170	30	200	93	1,225	17
1994	214	171	33	204	95	1,564	14
1995	231	187	31	218	94	1,668	14
1996	231	167	28	195	84	1,551	15
1997	276	201	35	236	86	1,767	16
1998	276	200	29	229	83	1,654	17
1999	379	280	46	326	86	2,235	17
2000	379	274	45	319	84	1,387	27
2001	406	305	52	357	88	1,472	28
2002	406	262	45	307	76	1,529	27
2003	469	265	94	359	77	1,495	31
2004	469	287	95	382	81	1,387	34
2005	350	191	90	281	80	1,471	21
2006	350	183	92	275	79	1,311	27
2007	350	203	76	280	80	1,505	23
2008	350	183	85	268	77	1,498	23
2009	260	147	53	200	77	1,339	19
2010	260	162	53	215	83	1,276	20
2011	235	144	45	189	80	1,393	17
2012	235	156	45	201	86	1,382	17

Table 2. Known moose mortalities, excluding controlled hunts, Upper Snake Region, 1982-present.

Year	Mortality agent						Total
	Native American Harvest	Illegal kill	Road kill	Natural	Train kill	Other	
1982	0	3	0	0	0	0	3
1983	0	6	4	0	0	2	12
1984	11	10	6	3	0	17	47
1985	6	12	13	1	6	9	47
1986	6	19	14	1	0	7	47
1987	6	14	14	7	2	8	51
1988	1	6	31	7	4	41	90
1989	2	2	10	1	0	9	24
1990	3	8	16	4	0	13	44
1991	1	10	12	6	4	22	55
1992	3	10	38	0	0	15	66
1993	1	8	7	0	0	4	20
1994	0	9	36	3	0	6	54
1995	2	3	15	2	0	7	29
1996	2	1	30	1	0	16	50
1997	1	7	27	9	0	5	49
1998	0	2	25	1	0	7	35
1999	2	4	26	5	0	3	40
2000	2	6	19	1	0	4	32
2001	0	3	11	1	0	9	24
2002	0	0	15	3	0	4	22
2003	0	2	14	3	0	0	19
2004	0	6	22	0	0	7	25
2005	0	1	27	5	0	6	39
2006	0	2	23	1	0	5	31
2007	0	1	1	9	0	2	13
2008	1	0	2	0	0	2	5
2009	0	1	0	3	0	3	7
2010	0	1	0	1	0	0	2
2011	0	3	0	1	0	16	20
2012	0	2	0	1	0	7	10

Table 3. Moose harvest and drawing odds by analysis area, Upper Snake Region, 2000-present.

Analysis area	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing odds (%)
			M	F				
50, 51,	2000	34	17	11	82	2.7	90	38
58, 63	2001	37	18	13	84	3.3	113	33
63A	2002	37	22	11	89	6.7	111	33
	2003	53	23	14	70	3.7	107	50
	2004	53	25	19	83	5.0	135	39
	2005	45	21	19	89	4.8	158	28
	2006	45	16	17	73	4.8	190	24
	2007	45	20	15	78	4.0	170	26
	2008	45	18	14	71	6.4	174	26
	2009	35	20	12	91	6.7	225	16
	2010	35	14	13	77	5.3	191	18
	2011	40	19	12	78	5.0	236	17
	2012	40	20	12	80	4.8	226	18
59, 59A	2000	20	19	0	95	4.8	110	18
	2001	22	19	0	86	2.6	88	25
	2002	22	20	0	91	6.7	124	18
	2003	25	20	5	100	5.0	113	22
	2004	25	19	5	96	3.1	102	25
	2005	20	12	3	75	4.5	131	15
	2006	20	14	5	95	2.3	85	24
	2007	20	13	4	85	4.4	109	18
	2008	20	15	4	95	6.1	74	27
	2009	5	3	0	60	11.0	39	13
	2010	5	4	4	80	9.0	30	17
	2011	5	5	0	100	11.8	45	11
	2012	5	5	0	100	4.8	48	10
60, 60A	2000	136	104	5	80	4.5	582	23
61, 62,	2001	144	119	13	92	4.2	651	22
62A	2002	144	94	9	72	7.2	616	23
	2003	174	89	32	70	5.9	605	29
	2004	174	103	33	78	5.2	516	34
	2005	120	63	29	77	5.4	532	23
	2006	120	66	30	80	5.2	448	27
	2007	120	73	22	79	5.4	531	23
	2008	120	59	29	73	5.7	479	25
	2009	80	50	13	79	6.5	408	20
	2010	80	49	12	76	7.3	379	21
	2011	65	45	8	82	6.5	420	15
	2012	65	49	9	89	5.5	384	17
64, 65,	2000	79	51	10	77	4.8	202	39
67	2001	74	55	9	86	3.8	175	42
	2002	74	41	8	66	6.8	217	34

Table 3. Continued.

Analysis area	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing odds (%)
			M	F				
	2003	78	48	16	82	8.7	184	42
	2004	78	47	14	78	6.2	230	34
	2005	65	36	14	77	5.5	205	32
	2006	65	31	17	74	5.7	198	33
	2007	65	40	15	85	6.8	236	28
	2008	65	38	13	78	7.1	256	25
	2009	65	30	14	68	7.5	275	24
	2010	65	38	15	82	8.6	295	22
	2011	60	30	17	78	6.6	296	20
	2012	61 ^a	31	18	80	7.1	291	21
66, 69	2000	110	83	19	93	5.3	403	27
	2001	129	94	17	86	5.2	445	29
	2002	129	85	17	79	6.8	461	28
	2003	139	81	29	79	5.3	486	29
	2004	139	92	26	85	5.3	404	34
	2005	100	59	25	84	6.6	445	22
	2006	100	56	23	79	5.8	390	26
	2007	100	56	20	76	5.7	459	22
	2008	100	53	25	78	7.1	345	29
	2009	75	44	13	76	7.3	392	19
	2010	75	53	11	85	5.4	295	25
	2011	65	45	5	77	7.7	396	16
	2012	65	51	6	88	6.2	433	15

^a Includes one super tag.

Table 4. Aerial survey of moose, Hunt Area 62, Upper Snake Region, 2000-2001.

	Observed	Estimated ($\pm 90\%$ CI)
Total moose	332	366 \pm 16
Cows	164	180 \pm 9
Bulls	98	109 \pm 8
Calves	70	77 \pm 5
Bulls:cows:calves	60:100:43	61:100:43

Table 5. Aerial survey of moose, Hunt Areas 60, 60A, 61, 62, Upper Snake Region.

Inclusive location	1990-1991		1991-1992	
	Bulls:cows:calves	Total	Bulls:cows:calves	Total
Middle to N Leigh Creek	67:100:83	15		0
Wiggleton Hollow to Johns Creek	56:100:56	19		7
N Fork Badger Creek to Bitch Crk	72:100:56	41		6
Bitch Creek to Conant Creek	7:100:68	49	56:100:67	20
Conant Creek to Fall River		14	27:100:55	20
Fall River Ridge to Cave Falls Rd	36:100:43	80		28
Cave Falls Rd to Fish Creek Rd		10	56:100:22	16
Fish Creek to Moose Creek		24		19
Warm River Hatchery to Survey Draw	17:100:67	11		5
Buffalo River		2		2
Macks Inn/Big Springs Henrys Lake Flat	42:100:52	59		19
Henrys Lake	22:100:56	16		19
Henrys Fork to Hatchery Butte west of Warm River	32:100:60	102		14
Humphrey to Spencer	73:100:55	25		14
Spencer to Rattlesnake Creek	25:100:75	24		23
Corral Creek to Spring Creek	5:100:47	29		7
West Camas Drainage		14		29
East Camas Drainage		9		4
Big Bend Ridge	14:100:105	88	22:100:122	68
Desert, east of Sand Creek		6		8
Desert, Red Rd to Sand Creek Rd ^a	100:100:100	85	65:100:41	50
Junipers and Hook of Sands ^a	118:100:44	103	33:100:67	18
Chokecherry Ridge and Second Sands ^a	69:100:45	63	72:100:36	48
Total		888		444

^a Moose counted in conjunction with helicopter deer survey, 18 December 1988.

SALMON REGION

GMUs 21, 21A, 27, 29, 30, 30A, 36A, 37A

Controlled Hunt Areas 21, 27, 29, 36A

Abstract

Three controlled hunts with 13 total tags for antlered moose occurred in Salmon Region during 2012. Hunt Area 21 was closed to hunting. Twelve of 13 hunters harvested moose (92% hunter success). Average antler spread was 38.2 inches; the 5-year running average was 37.5 inches. Interest in moose tags was typical of recent years; 124 applicants selected Salmon Region hunts as first choices (drawing odds = 11%).

Climatic Conditions

Rainfall during summer months in 2012 was slightly below average, with some cool, moist weather during spring and early summer. Vegetative growth appeared average. Winter conditions were relatively mild, with normal to above normal temperatures and below normal precipitation, at least at mid to lower elevations. In general, animals should have entered winter in above average body condition, and then encountered a mild winter, which should have produced high over-winter survival for adults. Snow-pack (as measured at higher elevations) was approximately 96% of average by late winter, 2013. Onset of spring weather and associated plant phenology was about normal in 2013. Water-year precipitation through June 2013 has been approximately 88% of average at lower elevations (Salmon weather station) and 87% of average at higher elevations (Moose Creek snotel site). Spring and early summer conditions in 2013 were cool and moist.

Background

Habitats in these GMUs range from riparian river bottoms to sagebrush grasslands on rolling foothills up through ponderosa pine and Douglas fir forests to lodgepole pine and spruce-fir forests at higher elevations. Willow shrub communities usually associated with moose habitat are not common. Portions of these GMUs contain extensive cliff and rock talus areas at both low and high elevations. Topography is moderately to very rugged. Game management units 21 and 21A are in one of the higher precipitation zones in Salmon Region, creating productive commercial forestlands. As a consequence, timber harvest is a dominant activity in at least the North Fork Salmon River drainage. Logging roads are common.

GMUs 21, 21A, 30, and 30A border areas in Montana where moose are common. Migrants from Montana may well have formed the initial nucleus for populations in GMUs bordering Montana. Cross-border movements are no doubt common in this area. No information exists on historical moose numbers other than an increase in moose sightings in recent decades. As a result, Hunt Area 21 (GMUs 21 and 21A) was initiated in 1990 with 3 tags. Similar increases in moose sightings resulted in establishment of Hunt Area 29 (GMUs 29 and 37A) in 1991 and Hunt Area 30 (GMUs 30 and 30A) in 1993. Hunt Area 30 was incorporated into Hunt Area 29 in 1999. Two new hunt areas were opened in 2005 with 1 tag each: 27 and 36A. GMU 58 (Upper Snake

Region) was added to Hunt Area 29 beginning in 2007. Hunt Area 21 was closed beginning in 2011 because of low hunter success in recent years (Figure 12).

Population Surveys

Because of dense cover, low moose densities, and solitary habits of moose, formal population surveys are generally ineffective in occupied moose habitat in Salmon Region. Incidental observations of moose are recorded during aerial surveys for other ungulates. During 2010-2011 surveys, observers counted 93 moose. Because most mule deer or elk surveys were cancelled in 2011-2012, no moose were observed. Mule deer and elk surveys were conducted in December, 2012 and February, 2013, respectively. Twenty-seven moose were observed during the elk survey: 7 cows, 1 calf, 18 bulls and 1 unknown. One additional moose was observed during a March 2013 mountain goat survey.

Harvest Characteristics

Harvest and hunter information was compiled from Big Game Mortality Reports, which hunters must complete within 10 days of harvest; antlers of males must be presented to a Department representative. Tag levels (Table 1) and season structure (Appendix A) were unchanged for Hunt Areas 27, 29, and 36A; Hunt Area 21 with 2 tags was closed 2011. Two tags were added in 2 new hunt areas in 2005 (Table 2); 1 tag each in areas 27 (all of GMU 27) and 36A (all of GMU 36A). Thirteen antlered-moose tags were allocated between 3 controlled hunts in Salmon Region for 2012. Twelve of 13 hunters harvested moose (92% success). Overall hunter success has increased over the long-term average of approximately 84%. Of 287 hunters since 1990, 228 (80%) have taken a moose (Table 1). Antler spread of moose harvested during the 2012 season ranged from 29 to 44.75 inches (mean = 39.2 in.). Since 1995, average spread ranged from 33.4 to 38.6 inches.

Two moose deaths were attributed to non-hunting mortality during the reporting period (Table 3). Non-hunting mortality ranged from 0 to 8 moose per year since 1982.

Habitat Conditions

Past logging operations in primary moose range of GMUs 21 and 21A generally enhanced moose habitat by encouraging forb and shrub production in cutover areas. However, positive impacts may eventually be counter-balanced by negative effects of increased road access and loss of mature, dense-canopy forest stands used by moose for winter cover. Timber harvest has declined in recent years and harvested areas are returning to pole stands with little forage value.

Capture and Translocation

No moose capture or translocation operations were conducted in Salmon Region during the reporting period (Table 4). Opportunities exist to expand moose populations in GMUs 36 and 36B via capture and translocation.

Table 1. Moose harvest and drawing odds, Salmon Region, 1990-present.

Year	Tags	Harvest			Hunter success (%)	First-choice applicants	Drawing odds (%)
		M	F	Total			
1990	3	2	0	2	67	12	25
1991	6	6	0	6	100	38	16
1992	6	6	0	6	100	32	19
1993	9	9	0	9	100	54	17
1994	9	8	0	8	89	54	17
1995	12	10	0	10	83	123	10
1996	12	11	0	11	92	82	15
1997	12	12	0	12	100	89	13
1998	12	11	0	11	92	92	13
1999	14	13	0	13	93	124	11
2000 ^a	14	11	0	11	79	80	18
2001 ^{a,b}	15	16	0	16	107	102	15
2002	14	12	0	12	86	76	18
2003	14	11	0	11	79	106	13
2004	14	11	0	11	79	93	15
2005 ^c	16	9	0	9	53	124	13
2006	16	13	0	13	81	119	13
2007	16	13	0	13	81	111	14
2008	16	11	0	11	69	113	14
2009	15	10	0	10	67	119	13
2010	16 ^d	13	0	13	81	116	14
2011	13	11	0	11	85	111	12
2012	13	12	0	12	92	124	11

^a One tag was deferred from 2000 until 2001 season because of wildfires.

^b Two hunters mistakenly harvested bulls in Hunt Area 29.

^c One hunter mistakenly harvested a bull in Hunt Area 29.

^d Super-tag holder hunted and harvested a bull (GMU 36A).

Table 2. Moose harvest and drawing odds by Hunt Area, Salmon Region, 1997-present.

Hunt Area	Year	Tags	Harvest		Hunter success (%)	Days/hunter	First-choice applicants	Drawing odds (%)
			M	F				
21	2000 ^a	4	2	0	67	4.0	10	40
	2001 ^a	5	4	0	80	16.3	15	33
	2002	4	2	0	50	10.5	15	27
	2003	4	3	0	75	9.0	10	40
	2004	4	3	0	75	7.0	9	44
	2005	4	1	0	25	16.0	11	36
	2006	4	2	0	50	12.5	9	44
	2007	4	2	0	50	6.0	4	100
	2008	4	1	0	25	11.0	6	67
	2009	2	0	0	0	0	0	0
2010	2	0	0	0	0	5	40	
27	2005	1	0	0	0		2	50
	2006	1	0	0	0		1	100
	2007	1	1	0	100	10.0	4	25
	2008	1	0	0	0		2	50
	2009	1	0	0	0		5	20
	2010	1	0	0	0		2	50
	2011	1	0	0	0		2	50
2012	1	0	0	0				
29	2000	10	9	0	90	4.9	70	14
	2001 ^b	10	12	0	100	6.7	87	11
	2002	10	10	0	100	7.9	61	16
	2003	10	8	0	80	6.3	96	10
	2004	10	8	0	80	7.0	84	12
	2005 ^c	10	8	0	73	4.0	108	9
	2006	10	10	0	100	6.4	91	11
	2007	10	9	0	90	5.1	87	11
	2008	10	10	0	100	5.7	97	10
	2009	11	9	0	82	11.0	99	11
	2010	11	11	0	100	4.7	99	11
	2011	11	10	0	91	8.1	102	11
2012	11	10	0	91	6.0	108	10	
30	1997	3	3	0	100	3.0	27	11
	1998 ^d	3	3	0	100	8.3	30	10
36A	2005	1	0	0	0		3	33
	2006	1	1	0	100	3.0	18	6
	2007	1	1	0	100	10.0	16	6
	2008	1	0	0	0		8	13
	2009	1	1	0	100	4.0	15	7
	2010	2	2	0	100	16.5	10	20
	2011	1	1	0	100	2.0	7	14
2012	1	1	0	100	2	16	6	

^a One tag was deferred from 2000 until 2001 season because of wildfires.

^b Two hunters mistakenly harvested bulls in Hunt Area 29.

^c One hunter mistakenly harvested a bull in Hunt Area 29.

^d Hunt Area 30 combined with Hunt Area 29 after 1998.

Table 3. Known moose mortalities, excluding controlled hunts, Salmon Region, 1982-present.

Year	Mortality agent					Total
	Native American harvest	Illegal kill	Road kill	Natural	Other	
1982	0	0	0	0	0	0
1983	0	0	0	0	0	0
1984	0	0	0	0	0	0
1985	0	0	0	0	0	0
1986	0	0	1	0	0	1
1987	0	0	0	1	0	1
1988	0	1	0	0	0	1
1989	0	0	0	0	0	0
1990	2	0	1	1	0	4
1991	6	0	0	0	0	6
1992	6	1	1	0	0	8
1993	0	1	0	1	0	2
1994	0	1	1	1	0	3
1995	0	0	0	2	0	2
1996	0	0	0	0	2	2
1997	0	1	1	1	0	3
1998	0	1	0	0	2	3
1999	0	0	1	0	1	2
2000	0	0	2	0	0	2
2001	0	2	2	0	0	4
2002	0	2	1	1	1	5
2003	0	0	3	1	0	4
2004	0	0	3	2	1	6
2005	0	1	0	1	1	3
2006	0	0	1	1	1	3
2007	0	0	0	1	0	1
2008	0	0	1	0	0	1
2009	0	1	3	0	0	4
2010	0	0	0	0	0	0
2011	0	0	0	0	0	0
2012	0	0	2	0	0	0

Table 4. Moose translocation, Salmon Region, February 1993.

Capture site	Release site	Adults		Calves		Total
		M	F	M	F	
GMUs 60, 60A, 62 in various locations	GMU 36: Valley Cr.	1	2	0	0	3
	GMU 36: Decker Flat	0	2	1	0	3
	GMU 36: Gold Cr.	0	2	0	0	2

APPENDIX A
IDAHO
2012 SEASON
MOOSE RULES

Moose, Bighorn Sheep and Mountain Goat

Controlled Hunt Seasons and Rules 2011 and 2012



Major changes are highlighted in yellow.



**Controlled Hunt
application
period:
April 1 - April 30.**

**Persons applying
for controlled
hunts must
submit tag and
application fees.**



2011 & 2012 Moose Hunting Seasons

Antler Restrictions

- Only moose with at least one antler longer than six inches may be taken in any season open for antlered moose only.
- Only moose without antlers or with antlers less than six inches long may be taken in any season which is open for antlerless moose only.

Mandatory Check and Report Requirements

Antlers must be presented at Idaho Fish and Game regional offices or official check point or to a conservation officer within 10 days of the date of the kill. Fish and Game headquarters office is not equipped to check in moose. In the Boise area, these animals can be checked at the IDFG Regional Office in Nampa (3101 S. Powerline Rd, 208-465-8465) between the hours of 8 a.m. and 5 p.m. or by appointment at the Garden City facility, 109 W. 44th St., 208-327-7099. Successful hunters must complete a big game mortality report, available at Fish and Game regional offices, from conservation officers, taxidermists

and meat processors within 10 days of the date of the kill. All hunters who have harvested either an antlered or antlerless moose must complete this report.

A hunter may authorize another person to comply with the above report requirements if that person complies with those requirements and possesses the necessary information to accurately complete the form.

Unsuccessful hunters must present or mail their unused tags to a Fish and Game office within 10 days after the close of the season for which the tag was valid. Cancelled tags will be returned to the hunter upon request. Failure to report may result in future ineligibility in moose drawings.

NOTE: Moose tags unfilled after the first drawing are available to any Idaho hunter during a second drawing. (See page 34) Hunters who have previously harvested a bull and/or a cow moose and not eligible for the first drawing **may apply** for and receive one of these tags in the second drawing or as a left-over tag if tags are still available.

2011 - 2012 Antlered Moose Controlled Hunts - 761 Tags							
Hunt No.	Controlled Hunt Area	Tags	Season Dates	Hunt No.	Controlled Hunt Area	Tags	Season Dates
3001	1-1	50	Sep 15-Dec 1	3025	7	5	Sep 15-Dec 1
3002	1-1	15	Oct 1-Oct 14	3026	7	5	Oct 1-Oct 14
3003	1-1	15	Nov 1-Nov 14	3027	8	8	Aug 30-Nov 23
3004	1-2	20	Sep 15-Dec 1	3028	8A	8	Aug 30-Nov 23
3005	1-2	5	Oct 1-Oct 14	3029	9	5	Sep 15-Dec 1
3006	1-2	10	Nov 1-Nov 14	3030	9	5	Oct 1-Oct 14
3007	1-3	10	Sep 15-Dec 1	3031	10-1	6	Aug 30-Nov 23
3008	1-4	30	Sep 15-Dec 1	3032	10-2	5	Aug 30-Nov 23
3009	1-4	5	Oct 1-Oct 14	3033	10-3	10	Aug 30-Nov 23
3010	1-4	10	Nov 1-Nov 14	3034	10-4	4	Aug 30-Nov 23
3011	2	20	Sep 15-Dec 1	3035	10-5	4	Aug 30-Nov 23
3012	2	10	Oct 1-Oct 14	3036	10-6	3	Aug 30-Nov 23
3013	2	10	Nov 1-Nov 14	3037	10A-1	5	Aug 30-Nov 23
3014	3	15	Sep 15-Dec 1	3038	10A-2	8	Aug 30-Nov 23
3015	3	5	Oct 1-Oct 14	3039	10A-3	3	Aug 30-Nov 23
3016	3	8	Nov 1-Nov 14	3040	10A-4	8	Aug 30-Nov 23
3017	4	15	Sep 15-Dec 1	3041	10A-5	5	Aug 30-Nov 23
3018	4	5	Oct 1-Oct 14	3042	12-1	2	Aug 30-Nov 23
3019	4	10	Nov 1-Nov 14	3043	12-2	10	Aug 30-Nov 23
3020	4A	5	Sep 15-Dec 1	3044	12-3	4	Aug 30-Nov 23
3021	5	15	Sep 15-Dec 1	3045	12-4	2	Aug 30-Nov 23
3022	6	15	Sep 15-Dec 1	3046	12-5	4	Aug 30-Nov 23
3023	6	5	Oct 1-Oct 14	3047	12-6	4	Aug 30-Nov 23
3024	6	5	Nov 1-Nov 14	3048	14-1	7	Aug 30-Nov 23

MOOSE

Hunt No.	Controlled Hunt Area	Tags	Season Dates
3049	14-2	4	Aug 30-Nov 23
3050	15	8	Aug 30-Nov 23
3051	16-1	2	Aug 30-Nov 23
3052	16-2	2	Aug 30-Nov 23
3053	16A	4	Aug 30-Nov 23
3054	17	5	Aug 30-Nov 23
3055	19	5	Aug 30-Nov 23
3056	20	5	Aug 30-Nov 23
3057	27	1	Aug 30-Nov 23
3058	29*	11	Aug 30-Nov 23
3059	36A	1	Aug 30-Nov 23
3060	44*	3	Aug 30 - Nov 23
3061	50	5	Aug 30-Nov 23
3062	51	5	Aug 30-Nov 23
3063	55*	5	Aug 30 - Nov 23
3064	59*	5	Aug 30-Nov 23
3065	60 ^a	15	Aug 30-Nov 23
3066	60A ^a	5	Aug 30-Nov 23
3067	61-1	10	Aug 30-Nov 23
3068	61-2	5	Aug 30-Nov 23
3069	61-3	10	Aug 30-Nov 23
3070	62	5	Aug 30-Nov 23
3071	62A	5	Aug 30-Nov 23
3072	63A ^{b,c}	15	Aug 30-Nov 23
3073	64	10	Aug 30-Nov 23
3074	65	10	Aug 30-Nov 23
3075	66-1	10	Aug 30-Nov 23
3076	66-2	10	Aug 30-Nov 23
3077	66A	10	Aug 30-Nov 23

Hunt No.	Controlled Hunt Area	Tags	Season Dates
3078	67-1	10	Aug 30-Nov 23
3079	67-2	10	Aug 30-Nov 23
3080	69-1	15	Aug 30-Nov 23
3081	69-2	15	Aug 30-Nov 23
3082	69-3*	5	Aug 30-Nov 23
3083	70	5	Aug 30-Nov 23
3084	71-1	5	Aug 30-Nov 23
3085	71-2	5	Aug 30-Nov 23
3086	72	5	Aug 30-Nov 23
3087	73*	5	Aug 30-Nov 23
3088	74	5	Aug 30-Nov 23
3089	75	5	Aug 30-Nov 23
3090	76-1	10	Aug 30-Nov 23
3091	76-2	7	Aug 30-Nov 23
3092	76-3	10	Aug 30-Nov 23
3093	77	5	Aug 30-Nov 23
3094	78	5	Aug 30-Nov 23

* See controlled hunt area descriptions. This hunt includes other units or parts of other units.
^a Short-range weapons only on Chester Wetlands WMA.
^b Short-range weapons only. Limited access.
^c Motorboat advised for game retrieval.



IDFG photo by Dale Towell

↓ **2011 - 2012 Antlerless Moose Controlled Hunts - 163 Tags**

Hunt No.	Controlled Hunt Area.	Tags	Season Dates	Hunt No.	Controlled Hunt Area.	Tags	Season Dates
3095	1-1	10	Oct 15-Dec 1	3108	67-1	5	Oct 15-Nov 23
3096	2	35	Oct 15-Dec 1	3109	67-2	5	Oct 15-Nov 23
3097	3	5	Oct 15-Dec 1	3110	69-1	5	Oct 15-Nov 23
3098	5	5	Oct 15-Dec 1	3111	69-2	5	Oct 15-Nov 23
3099	8	4	Oct 15-Nov 23	3112	71-1	5	Oct 15-Nov 23
3100	8A	4	Oct 15-Nov 23	3113	71-2	5	Oct 15-Nov 23
3101	50	5	Oct 15-Nov 23	3114	75	5	Oct 15-Nov 23
3102	60 ^a	5	Oct 15-Nov 23	3115	76-1	5	Oct 15-Nov 23
3103	60A ^b	5	Oct 15-Nov 23	3116	76-2	5	Oct 15-Nov 23
3104	63A ^{b,c}	10	Oct 15-Nov 23	3117	76-3	5	Oct 15-Nov 23
3105	64	5	Oct 15-Nov 23	3118	77	5	Oct 15-Nov 23
3106	65	5	Oct 15-Nov 23	3119	78	5	Oct 15-Nov 23
3107	66A	5	Oct 15-Nov 23				

^a Short-range weapons only on Chester Wetlands WMA.
^b Short-range weapons only. Limited access.
^c Motorboat advised for game retrieval.

HUNT AREA DESCRIPTIONS

Hunt Area 1-1 — That portion of Unit 1 within the Priest River drainage, and those portions of the Pend Oreille and Salmon River drainages downstream from the Priest River drainage.

Hunt Area 1-2 — That portion of Unit 1 within the following boundaries: beginning on U.S. Highway 95 bridge across the Pend Oreille River at Sandpoint, then northward along Highway 95 to the Kootenai River at Bonner's Ferry, then northwesterly along the Kootenai River to the U.S. border, then west along the U.S. border to the Priest River-Kootenai River divide, then south along the Priest River-Pack River divide to Flat Top Mountain, then south along the divide separating the Priest River drainage and the Pend Oreille drainage to Priest River, then east along the Pend Oreille River to the point of beginning. **Except Myrtle Creek Game Preserve - Closed.**

Hunt Area 1-3 — That portion of Unit 1 north and east of the Kootenai River.

Hunt Area 1-4 — That portion of Unit 1 south of the Kootenai River and east of U.S. Highway 95. **Except The David Thompson Game Preserve - Closed.**

Hunt Area 2 — All of Unit 2.

Hunt Area 3 — All of Unit 3.

Hunt Area 4 — All of Unit 4.

Hunt Area 4A — All of Unit 4A.

Hunt Area 5 — All of Unit 5.

Hunt Area 6 — All of Unit 6.

Hunt Area 7 — All of Unit 7.

Hunt Area 8 — All of Unit 8.

Hunt Area 8A — All of Unit 8A.

Hunt Area 9 — All of Unit 9.

Hunt Area 10-1 — That portion of Unit 10 within the Cayuse Creek drainage.

Hunt Area 10-2 — That portion of Unit 10 on the north side of the Kelly Creek drainage upstream from, but excluding, the Moose Creek drainage, and that portion on the south side of the Kelly Creek drainage upstream from, but excluding, the Cayuse Creek drainage.

Hunt Area 10-3 — That portion of Unit 10 on the north side of the Kelly Creek drainage upstream from its mouth to and including the Moose Creek drainage, and the North Fork of the Clearwater River drainage upstream from the mouth of Kelly Creek.

Hunt Area 10-4 — That portion of Unit 10 within the Fourth of July Creek drainage, that portion on the south side of the North Fork of the Clearwater River from the mouth of Fourth of July Creek upstream to the mouth of Kelly Creek, and the south side of the Kelly Creek drainage from its mouth upstream to, but excluding, the Cayuse Creek drainage.

Hunt Area 10-5 — That portion of Unit 10 within the Weitas Creek drainage (a tributary of the upper North Fork of the Clearwater River), and the drainages on the southwest side of the North Fork of the Clearwater River from the Weitas Creek drainage to, but excluding, the Fourth of July Creek drainage.

Hunt Area 10-6 — That portion of Unit 10 on the north side of the North Fork of the Clearwater River drainage downstream from the mouth of Kelly Creek.

Hunt Area 10A-1 — That portion of Unit 10A within the following boundary: Beginning at the junction of the Unit 10A boundary with Forest Service Road 250 along the North Fork of the Clearwater River, then southwest along Forest Service Road 250 to Forest Service Road 669, then west and south along Forest Service Road 669 to Highway 11 at Pierce, then south on Highway 11 to Forest Service Road 100, then south on Forest Service Road 100 to the Clearwater National Forest boundary, then south along the Clearwater National Forest boundary to the Unit 10A boundary, then north along the Unit 10A boundary to the point of beginning.

Hunt Area 10A-2 — That portion of Unit 10A within the following boundary: Beginning at the junction of Unit 10A boundary with Forest Service Road 247, then south on Forest Service Road 247 to Forest Service Road 251, then south on Forest Service Road 251 to Forest Service Road 246, then southwest on Forest Service Road 246 to State Highway 11 at Headquarters, then south on Highway 11 to Forest Service Road 669 at Pierce, then northeast on Forest Service Road 669 to Forest Service Road 250, then northeast on Forest Service Road 250 to the Unit 10A boundary, then north and east along the Unit 10A boundary to the point of beginning.

Hunt Area 10A-3 — That portion of Unit 10A within the following boundary: Beginning at the Grandad Bridge on the Unit 10A boundary, then south and east along the Silver Creek-Casey Creek Road to Forest Service Road 247, then south on Forest Service Road 247 to Forest Service Road 246 at Headquarters, then northeast on Forest Service Road 246 to Forest Service Road 251, then north on Forest Service Road 251 to Forest Service Road 247, then north on Forest Service Road 247 to the Unit 10A boundary at the North Fork of the Clearwater River, then west on the Unit 10A boundary to the point of beginning.

Hunt Area 10A-4 — That portion of Unit 10A north of Forest Service Road 1705 from Elk River to Grandad Bridge and north and west of Dworshak Reservoir and the Little North Fork of the Clearwater River.

Hunt Area 10A-5 — That portion of Unit 10A south of Forest Service Road 1705 from Elk River to Grandad Bridge and north and west of Dworshak Reservoir.

Hunt Area 12-1 — That portion of Unit 12 north of the Lochsa River from and including the Lost Creek drainage upstream to, but excluding the Crooked Fork drainage.

Hunt Area 12-2 — That portion of Unit 12 within the Crooked Fork drainage and north of White Sand Creek upstream to and including the Storm Creek drainage.

Hunt Area 12-3 — That portion of Unit 12 south of the Lochsa River from and including the Old Man Creek drainage upstream to and including the Mocus Creek drainage.

Hunt Area 12-4 — That portion of Unit 12 south of the Lochsa River from, but excluding, the Mocus Creek drainage, upstream to and including the Cliff Creek drainage.

Hunt Area 12-5 — That portion of Unit 12 within the Walton Creek drainage, that portion on the south side of White Sand Creek upstream to the mouth of Storm Creek, and all of White Sand Creek drainage upstream from, but excluding, the Storm Creek drainage.

Hunt Area 12-6 — That portion of Unit 12 north of the Middle Fork of the Clearwater River from the Smith Creek Road (Forest Service Road 101) upstream to the mouth of the Lochsa River, that portion

on the north side of the Lochsa River upstream to, but excluding, the Lost Creek drainage, and that portion on the south side of the Lochsa River from its mouth upstream to, but excluding, the Old Man Creek drainage.

Hunt Area 14-1 — That portion of Unit 14 north of the following boundary: Beginning on the Unit 14 west boundary on the Slate Creek Road (Forest Service Road 354), then east on the Slate Creek Road to Forest Service Road 221, then north on Forest Service Road 221 to the Unit 14 east boundary.

Hunt Area 14-2 — That portion of Unit 14 south of the following boundary: Beginning on the Unit 14 west boundary on the Slate Creek Road (Forest Service Road 354), then east on the Slate Creek Road to Forest Service Road 221, then north on Forest Service Road 221 to the Unit 14 east boundary.

Hunt Area 15 — All of Unit 15.

Hunt Area 16-1 — That portion of Unit 16 north and west of the Hamby Creek Road (Forest Service Road 651), and that portion south and west of the Selway River from its mouth upstream to the Hamby Creek Road.

Hunt Area 16-2 — That portion of Unit 16 south and east of Hamby Creek Road (Forest Service Road 651), and that portion north and east of the Selway River from its mouth upstream to Fog Mountain Road (Forest Service Road 319).

Hunt Area 16A — All of Unit 16A.

Hunt Area 17 — All of Unit 17.

Hunt Area 19 — All of Unit 19.

Hunt Area 20 — All of Unit 20.

Hunt Area 27 — All of Unit 27.

Hunt Area 29 — All of Units 29, 30, 30A, 37A, and 58.

Hunt Area 36A — All of Unit 36A.

Hunt Area 44 — That portion of Unit 44 east of the Fairfield-Couch Summit-Five Points Road and all of Units 48 and 49.

Hunt Area 50 — All of Unit 50.

Hunt Area 51 — All of Unit 51.

Hunt Area 55 — All of Units 55, 56 and 57.

Hunt Area 59 — All of Units 59 and 59A.

Hunt Area 60 — All of Unit 60.

Hunt Area 60A — That portion of Unit 60A south of the following line: from the intersection of 2300 N. (Egin-Hamer Road) and I-15 east to 3100 E. south to 2200 N. east to 1900 E. (Red Road) north to 600 N. (North Parker Road) east to US Hwy 20 business loop and north and then east on US Hwy business loop (Yellowstone Hwy) to US Hwy 20.

Hunt Area 61-1 — That portion of Unit 61 west of East Dry Creek.

Hunt Area 61-2 — That portion of Unit 61 east of East Dry Creek and west of U.S. Highway 191-20 and south and west of State Highway 87.

Hunt Area 61-3 — That portion of Unit 61 north of State Highway 87 and that portion east of U.S. Highway 191-20 except that portion enclosed by the Big Springs Loop Road and U.S. Highway 191-20.

Hunt Area 62 — All of Unit 62.

Hunt Area 62A — All of Unit 62A.

Hunt Area 63A — All of Unit 63A.

Hunt Area 64 — All of Unit 64.

Hunt Area 65 — All of Unit 65.

Hunt Area 66-1 — That portion of Unit 66 north of main Bear Creek except the Pritchard and Garden Creek drainages.

Hunt Area 66-2 — That portion of Unit 66 south of main Bear Creek.

Hunt Area 66A — All of Unit 66A.

Hunt Area 67-1 — That portion of Unit 67 north and west of Highway 31 and north of Highway 26.

Hunt Area 67-2 — That portion of Unit 67 south and east of Highway 31 and that portion of Unit 67 south of Highway 26.

Hunt Area 69-1 — That portion of Unit 69 west of the Grays Lake-Long Valley-Bone-Iona Road.

Hunt Area 69-2 — That portion of Unit 69 east of the Grays Lake-Long Valley-Bone-Iona Road **except** the Antelope and Granite Creek drainages.

Hunt Area 69-3 — That portion of Unit 69 within the Antelope and Granite Creek drainages, and that portion of Unit 66 within the Pritchard and Garden Creek drainages.

Hunt Area 70 — All of Unit 70.

Hunt Area 71-1 — That portion of Unit 71 located in Bannock and Bingham counties.

Hunt Area 71-2 — That portion of Unit 71 located in Caribou County.

Hunt Area 72 — All of Unit 72.

Hunt Area 73 — All of Units 73 and 73A.

Hunt Area 74 — All of Unit 74.

Hunt Area 75 — All of Unit 75.

Hunt Area 76-1 — That portion of Unit 76 within the following boundary: Beginning at Soda Springs on State Highway 34, then northeast to the Lanes Creek Road at Wayan, then south along the Lanes Creek-Diamond Creek Road to Timber Creek Road, then northeast along Timber Creek-Smoky Canyon-Stump Creek Road to the Idaho-Wyoming state line, then south along the state line to the Crow Creek Road, then southwest along Crow Creek-Wells Canyon-Georgetown Canyon Road to U.S. 30, then north along U.S. Highway 30 to Soda Springs, the point of beginning.

Hunt Area 76-2 — That portion of Unit 76 south of the Georgetown-Wells Canyon-Crow Creek Road.

Hunt Area 76-3 — That portion of Unit 76 north and east of the following boundary: Beginning at the Idaho-Wyoming state line, then west along the Stump Creek-Smoky Canyon-Timber Creek Road to the Diamond Creek Road, then north along the Diamond Creek-Lanes Creek Road to State Highway 34 at Wayan.

Hunt Area 77 — All of Unit 77.

Hunt Area 78 — All of Unit 78.



IDFG photo by Dale Towe III


Full text descriptions of hunt unit boundaries can be seen in the Idaho Administrative Procedures Act (<http://adm.idaho.gov/adminrules/rules/idapa13/>) or Big Game Rules brochure.

2011 - 2012 ANTLERED MOOSE HUNTS

Online users: Click on this link to view this map on your computer

<http://fishandgame.idaho.gov/ifwis/huntplanner/maps>

MOOSE



<http://fishandgame.idaho.gov>


9

Full text descriptions of hunt unit boundaries can be seen in the Idaho Administrative Procedures Act (<http://adm.idaho.gov/adminrules/rules/idapa13/>) or Big Game Rules brochure.

MOOSE

2011 - 2012 ANTLERED MOOSE HUNTS

Online users: Click on this link to view this map on your computer
<http://fishandgame.idaho.gov/ifwis/huntplanner/maps>




Full text descriptions of hunt unit boundaries can be seen in the Idaho Administrative Procedures Act (<http://adm.idaho.gov/adminrules/rules/idapa13/>) or Big Game Rules brochure.

2011 - 2012 ANTLERED MOOSE HUNTS

Online users: Click on this link to view this map on your computer
<http://fishandgame.idaho.gov/ifwis/huntplanner/maps>

MOOSE



<http://fishandgame.idaho.gov>

11


Full text descriptions of hunt unit boundaries can be seen in the Idaho Administrative Procedures Act (<http://adm.idaho.gov/adminrules/rules/idapa13/>) or Big Game Rules brochure.

MOOSE

2011 - 2012 ANTLERED MOOSE HUNTS

Online users: Click on this link to view this map on your computer

<http://fishandgame.idaho.gov/ifwis/huntplanner/maps>




Full text descriptions of hunt unit boundaries can be seen in the Idaho Administrative Procedures Act (<http://adm.idaho.gov/adminrules/rules/idapa13/>) or Big Game Rules brochure.

2011 - 2012 ANTLERLESS MOOSE HUNTS

Online users: Click on this link to view this map on your computer

<http://fishandgame.idaho.gov/ffwis/huntplanner/maps>



<http://fishandgame.idaho.gov>

13


MOOSE

Full text descriptions of hunt unit boundaries can be seen in the Idaho Administrative Procedures Act (<http://adm.idaho.gov/adminrules/rules/idapa13/>) or Big Game Rules brochure.

2011 - 2012 ANTLERLESS MOOSE HUNTS

Online users: Click on this link to view this map on your computer

<http://fishandgame.idaho.gov/fwis/huntplanner/maps>



Submitted by:

Wayne Wakkinen
Regional Wildlife Manager

George Pauley
Regional Wildlife Manager

Craig White
Regional Wildlife Manager

Regan Berkley
Regional Wildlife Manager


Randy Smith
Regional Wildlife Manager

Martha Wackenhut
Regional Wildlife Manager


Daryl Meints
Regional Wildlife Manager

Greg Painter
Regional Wildlife Manager

Approved by: IDAHO DEPARTMENT OF FISH AND GAME



Brad Compton, Asst. Chief
Bureau of Wildlife



Jeff Gould, Chief
Bureau of Wildlife